

Dalton Transactions

An international journal of inorganic chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: L. Cronin, C. Lin, M. Hutin, C. Busche, N. L. Bell and D. Long, *Dalton Trans.*, 2021, DOI: 10.1039/D0DT04149A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

Elucidating the Paramagnetic Interactions of an Inorganic-Organic Hybrid Radical-Functionalized Mn-Anderson Cluster

Chang-Gen Lin,^{a,b} Marie Hutin,^a Christoph Busche,^a Nicola Bell,^a De-Liang Long,^{a*} and Leroy Cronin^{a*}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

A family of six polyoxometalate-based magnetic compounds were synthesized by anchoring N-oxide type TEMPO radicals onto the Anderson type polyoxometalate cluster. The complexes were structurally characterised by single crystal X-ray diffraction and the intramolecular paramagnetic interactions between TEMPO radicals and Mn ions of the resulting hybrids were investigated in detail by electron paramagnetic resonance and the Evans NMR method.


Polyoxometalates (POMs) represent a vast class of negatively charged metal-oxo clusters¹ applicable to catalysis,² electronics,^{3,4} and magnetism.⁵ In particular, it has been reported that a highly redox stable POM, [PMo₁₂O₄₀(VO)₂]ⁿ⁻, shows promise for molecular spin-qubit quantum computing, with its molecular magnetism tuneable through electrical manipulation.⁶ Besides the fascinating redox properties of POMs, their enormous diversity in both size and structure makes them ideal building blocks to accommodate magnetic ions at specific sites, thus leading to magnetic clusters with definite topology and tuneable coordination environment.^{7,8}

Although big advances have been made on POM-based single molecular magnets, there currently is few examples of such systems based on covalently modified POM clusters. Hence in this context, we envision that it might be interesting to construct POM based magnetic molecules by anchoring stable organic radicals onto the metal-oxo cluster, and investigate the intramolecular magnetic interactions. Here we choose the Mn-Anderson cluster,⁹ [MnMo₆O₂₄{(CH₂)₃CNH₂}₃]³⁻, as the platform and covalently attach 2,2,6,6-tetramethyl-piperidine-1-oxyl (TEMPO) radicals on both sides of it. The magnetic properties of the resulting organic-inorganic hybrids were preliminarily monitored by electron paramagnetic resonance (EPR), and the interactions between the TEMPO radicals and the center Mn^{III}

ion were investigated systematically using the Evans NMR method, which has rarely been applied in POM chemistry.

The Evans method, first developed in 1959,¹⁰ has become a standard method to measure the magnetic susceptibility and unpaired electrons of inorganic complexes by using a NMR spectrometer.¹¹⁻¹³ The Evans NMR is performed in a diamagnetic solvent, and relies on the effect of the dissolved paramagnetic sample on the chemical shift of a reference compound, usually the solvent.^{14,15} The data is readily acquired by collecting the ¹H NMR spectrum of the paramagnetic sample that contains a capillary insert of pure solvent as reference.

The general synthetic procedure for the TEMPO-functionalized Mn-Anderson cluster is depicted in Scheme 1. It contains three steps: first, the carboxylic acid modification of Mn-Anderson cluster using glutaric anhydride; second, N-hydroxysuccinimide (NHS) activation; and third, amidation reaction using 4-amino-TEMPO. The biradical POM hybrid, [(n-C₄H₉)₄N]₃[MnMo₆O₂₄{(CH₂)₃CNHCO(CH₂)₃CONHC₉H₁₇NO}₂], is designated as **1-Mn**. Similarly, an analogue compound **2-Mn**, [(n-C₄H₉)₄N]₃[MnMo₆O₂₄{(CH₂)₃CNHCO(CH₂)₂CONHC₉H₁₇NO}₂], with a shorter alkyl chain, and an Al^{III} centered Anderson hybrid isostructure **3-Al**, [(n-C₄H₉)₄N]₃[AlMo₆O₂₄{(CH₂)₃CCH₂O-CO(CH₂)₂CONHC₉H₁₇NO}₂], were prepared for comparisons (See SI for details).



Scheme 1 General synthetic procedure of **1-Mn**. Colour code: Mo, Teal; Mn, Violet; and O, Red.

The biradical hybrids were characterized by FT-IR, ESI-MS, elemental analysis and single-crystal X-ray diffraction. Taking **1-Mn** as an example, the FT-IR spectrum shows typical bands for

^aWestCHEM, School of Chemistry, The University of Glasgow, Glasgow G12 8QQ, UK. E-mail: deliang.long@glasgow.ac.uk; lee.cronin@glasgow.ac.uk;

^bState Key Laboratory of Chemical Resource Engineering, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing 100029, P. R. China.

Electronic Supplementary Information (ESI) available: [detailed synthetic procedure, ESI-MS spectrum, EPR spectra, Evans NMR, and X-Ray crystallographic data]. See DOI: 10.1039/x0xx00000x



an Anderson cluster at 940 - 890 cm^{-1} (Mo=O) and 650 cm^{-1} (Mo-O-Mo), while the vibration of the amide bonds can be found at 1645 cm^{-1} . The molecular ion peak of **1-Mn** in ESI-MS spectrum is attributed to $m/z = 2174.40$ ($\{[\mathbf{1-Mn}]\text{-TBA}\}^+$). The X-ray structure of **1-Mn** is shown in Fig. 1. It crystallizes in a triclinic system with P-1 space group. Two unique half molecules (α and β) found in the asymmetrical unit. The Mn atoms in each half molecule locate on an inversion centre that is to generate the whole molecule. α and β differ in the conformation of TEMPO with the N-O lengths and related C-N-C angles very close to each other (Fig. 1A). Though slightly different, these geometrical parameters are all within the range of six-membered ring nitroxide.¹⁶ In crystal packing mode, each α molecule is surrounded by four β molecules at the ac plane, giving rise to a radical layer which further stacks along the b axis (Fig. 1B). No intermolecular contacts of TEMPO radicals are observed within the packing layer, or between layers. Any two N-O groups in the crystal packing are distant more than 5 Å.




Fig. 1 (A) The single-crystal structure of **1-Mn**, and (B) the crystal packing mode of **1-Mn** view along a axis: α molecule in red and β molecule in blue. Colour code: Mo, Teal; Mn, Violet; and O, Red; N, blue; and C, grey. H atoms, TBA counter cations, and solvent molecules are omitted for clarity.

The magnetic properties of these organic-inorganic biradicals were first investigated by EPR spectroscopy. As shown in Fig. 2, all the compounds exhibit the typical three-line signal of TEMPO due to the hyperfine splitting of the radical electrons with the nitrogen nucleus ($I = 1$),¹⁷ which indicates the absence of spin coupling between the two TEMPO groups within the biradical hybrid. This behaviour is expected since the N...N distance of the two TEMPO radicals within a molecule is too far (*ca.* 25 Å) to allow for strong J -coupling between the unpaired electrons.^{18,19} High-spin manganese(III) ($d4$, $S = 2$) is archetypical of non-Kramers ions resulting in Mn^{III} centres, which are "EPR-silent" due to a zero-field-splitting with an energy splitting higher than the microwave frequency used (X-Band).^{20,21} Possible signals resulting from the excited state could not be detected, due to the strong signal of the TEMPO radical.

However, comparison of the EPR spectra of **1-Mn** and **2-Mn** with that of **3-Al** allow us to conclude that the paramagnetic Mn^{III} has no effect on TEMPO radicals. To further confirm this, the Evans NMR method was adopted to explore the magnetic properties of these biradical hybrids.




Fig. 2 The EPR spectra of **1-Mn**, **2-Mn** and **3-Al** performed in acetonitrile solutions (1×10^{-4} M) at room temperature.

Even though the Evans NMR has been proved to be an effective method to study the solution state magnetic susceptibility of coordination complexes, the attempts to extend it in metal-oxo clusters, including the organically modified ones, have to our knowledge not been explored. Thus, in our first trial the paramagnetic properties of Mn-Anderson cluster were examined (See SI for experimental details). The calculation of the magnetic moment is given by Equation 1:


$$\mu_{exp} = 2.828 \sqrt{\frac{3T}{4\pi \cdot 10^6} \cdot \left[\left(\frac{\Delta_{ppm}}{c} \right)^{para} - \left(\frac{\Delta_{ppm}}{c} \right)^{dia} \right]} \quad (1)$$

where Δ_{ppm} is the chemical shift of DMSO- d_6 in ppm, c is the molar concentration of the solute (mol/mL), T is the probe temperature (K), and the *para* and *dia* superscripts stand for the paramagnetic and diamagnetic contributions, respectively. According to Piguet's work,¹⁵ the diamagnetic contribution of large paramagnetic assembly, here in our case Mn-Anderson, can be eliminated by deducting the diamagnetic contribution of an analogous assembly, where the paramagnetic metal ion is replaced by a proper diamagnetic one. Therefore, the Al^{III} centered Anderson cluster $[(n\text{-C}_4\text{H}_9)_4\text{N}]_3[\text{AlMo}_6\text{O}_{24}\{(\text{CH}_2)_3\text{C-CH}_2\text{OH}\}_2]$ (**3a**) and $[(n\text{-C}_4\text{H}_9)_4\text{N}]_3[\text{AlMo}_6\text{O}_{24}\{(\text{CH}_2)_3\text{CCH}_2\text{OCO-}(\text{CH}_2)_2\text{COOH}\}_2]$ (**3b**) were used. However, neither of these two Al^{III} analogues produce shifts in solvent chemical shift, thus the diamagnetic contribution can be taken as zero (Fig. S6). Thereafter, the magnetic moment (μ_{exp}) of Mn-Anderson is measured to be 4.69, which coincides with the theoretic result (4.90) obtained by Equation 2:

$$\mu_{theo} = \sqrt{n(n+2)} \quad (2)$$

where n is the number of unpaired electrons.

The implementation of the Evans method in Mn-Anderson inspired us to go further for the biradical hybrids. As shown in Fig. 3, the solvent chemical shifts in both **1-Mn** and **3-Al** are



observed. The corresponding magnetic moments are thus calculated based on Equation 1, and the results are listed in Table 1. The μ_{exp} value of **1-Mn** is found to be very close to μ_{theo} (low spin), suggesting the Mn^{III} might adopt a low spin state, that is, two unpaired electrons at the outer 3d orbital.




Fig. 3 The Evans NMR spectra of **1-Mn** and **3-Al**. Square and round dots indicate the solvent peaks of H₂O and DMSO, respectively. The chemical shifts of these solvents are labelled in graph.

However, this finding is contradictory to that suggested by the X-ray diffraction, as all the Mn-O bond lengths of **1-Mn** are similar to those of previously reported Mn-Anderson POMs (Fig. S1), which strongly indicates the Mn^{III} ion in biradical hybrid is still at its high spin state.⁹ Similar phenomena are also observed in the case of **2-Mn**. As for **3-Al**, μ_{calc} is much smaller than μ_{theo} , which implies a weak antiferromagnetic coupling of the magnetic centres.

As such, in order to obtain the magnetic moment of Mn^{III} in **1-Mn**, we hypothesize that the paramagnetic contribution of TEMPO may be eliminated by using **3-Al** as a reference compound, since these biradical hybrids are similar in both structure and molecular mass.¹⁵ To prove this, a modified equation (Equation 3) is used to calculate the magnetic moment of the Mn^{III} ion in **1-Mn**.

$$\mu_{\text{exp}}^{\text{Mn}} = 2.828 \sqrt{\frac{3 \cdot T}{4\pi \cdot 10^6} \cdot \left[\left(\frac{\Delta_{\text{ppm}}}{c} \right)^{1\text{Mn}} - \left(\frac{\Delta_{\text{ppm}}}{c} \right)^{3\text{Al}} \right]} \quad (3)$$

Table 1 Magnetic moments calculated by the Evans NMR method.

	μ_{exp}	μ_{theo} (high)	μ_{theo} (low)
Mn-Anderson ⁹	4.69 ± 0.04	4.90 (4)	-
1-Mn	5.22 ± 0.13	6.93 (6)	4.90 (4)
2-Mn	5.16 ± 0.15	6.93 (6)	4.90 (4)
3-Al	2.23 ± 0.09	2.83 (2)	-
Mn ^{III} in 1-Mn	4.72 ± 0.14	4.90 (4)	2.83 (2)
Mn ^{III} in 2-Mn	4.65 ± 0.16	4.90 (4)	2.83 (2)

The last two columns are the theoretic magnetic moments with the number of unpaired electrons given in bracket.

As listed in Table 1, the μ_{calc} of Mn^{III} in **1-Mn** is found to be 4.72, which is very close to that of Mn-Anderson (4.69). This result is in agreement with the EPR (non-Kramers ion) and X-ray data and indicates a high spin state of Mn^{III}. In the case of **2-Mn**, the μ_{exp} of Mn^{III} is calculated as 4.65, which further validates that there is no spin coupling between Mn^{III} and the TEMPO radicals.

Conclusions

In conclusion, we have represented here an example of magnetic molecules that are prepared by covalently anchoring TEMPO radicals onto the Anderson clusters. The resulting organic-inorganic biradicals have been fully characterized by various techniques such as ESI-MS and single-crystal X-ray diffraction. The solution state magnetic interactions between TEMPO radicals and the Mn^{III} ions have been investigated by EPR spectroscopy and Evans NMR. Though it seems there might exist some spin coupling, close investigation reveals that such 'false' interaction is due to the magnetic susceptibility calculation of TEMPO radicals using the Evans method, which turns out to be unsuitable. Therefore, the paramagnetic contribution of TEMPO in such biradical hybrids should be taken into account to get the actual spin state of Mn^{III}.

Conflicts of interest


There are no conflicts to declare.

Acknowledgements

This research was supported by EPSRC (grant nos. EP/J015156/1, EP/L023652/1, EP/I033459/1, EP/K023004/1) and ERC for an Advanced Grant (670467 SMART-POM). C.-G.L. thanks the financial support from China Scholarship Council and the National Natural Science Foundation of China (21901016).

Notes and references

- a) D.-L. Long, R. Tsunashima and L. Cronin, *Angew. Chem. Int. Ed.*, 2010, **49**, 1736; b) A. Proust, B. Matt, R. Villanneau, G. Guillemot, P. Gouzerh and G. Izzet, *Chem. Soc. Rev.*, 2012, **41**, 7605; c) A. V. Anyushin, A. Kondinski and T. N. Parac-Vogt, *Chem. Soc. Rev.*, 2020, **49**, 382; d) N. I. Gumerova, and A. Rompel, *Chem. Soc. Rev.*, 2020, **49**, 7568.
- S.-S. Wang and G.-Y. Yang, *Chem. Rev.*, 2015, **115**, 4893.
- C. Busche, L. Vila-Nadal, J. Yan, H. N. Miras, D.-L. Long, V. P. Georgiev, A. Asenov, R. H. Pedersen, N. Gadegaard, M. M. Mirza, D. J. Paul, J. M. Poblet and L. Cronin, *Nature*, 2014, **515**, 545.
- C. Kato, R. Machida, R. Maruyama, R. Tsunashima, X.-M. Ren, M. Kurmoo, K. Inoue and S. Nishihara, *Angew. Chem. Int. Ed.*, 2018, **57**, 13429.
- J. M. Clemente-Juan, E. Coronado and A. Gaita-Ariño, *Chem. Soc. Rev.*, 2012, **41**, 7464.
- J. Lehmann, A. Gaita-Ariño, E. Coronado and D. Loss, *Nat. Nanotech.*, 2007, **2**, 312.
- C. Ritchie, A. Ferguson, H. Nojiri, H. N. Miras, Y.-F. Song, D.-L. Long, E. Burkholder, M. Murrie, P. Kögerler, E. K. Brechin and L. Cronin, *Angew. Chem. Int. Ed.*, 2008, **47**, 5609.



COMMUNICATION

Journal Name

- 8 S. Cardona-Serra, J. M. Clemente-Juan, E. Coronado, A. Gaita-Ariño, A. Camón, M. Evangelisti, F. Luis, M. J. Martínez-Pérez and J. Sesé, *J. Am. Chem. Soc.*, 2012, **134**, 14982.
- 9 P. R. Marcoux, B. Hasenknopf, J. Vaissermann and P. Gouzerh, *Eur. J. Inorg. Chem.*, 2003, 2406.
- 10 D. F. Evans, *J. Chem. Soc.*, 1959, 2003.
- 11 A. A. Pavlov, G. L. Denisov, M. A. Kiskin, Y. V. Nelyubina and V. V. Novikov, *Inorg. Chem.*, 2017, **56**, 14759.
- 12 P. Lommens, P. Tack, L. V. Elst, I. V. Driessche, L. Vincze and D. Sinnaeve, *Dalton Trans.*, 2018, **47**, 3755.
- 13 T. Tezgerevska, E. Rousset, R. W. Gable, G. N. L. Jameson, E. C. Sañudo, A. Starikova and C. Boskovic, *Dalton Trans.*, 2019, **48**, 11674.
- 14 D. H. Grant, *J. Chem. Educ.*, 1995, **72**, 39.
- 15 C. Piguet, *J. Chem. Educ.*, 1997, **74**, 815.
- 16 H. Karoui, F. L. Moigne, O. Ouari and P. Tordo, in *Stable Radicals* (Editor, R. G. Hicks), 2010, Wiley, West Sussex, U.K.
- 17 A. L. Barra, L. C. Brunel and J. B. Robert, *Chem. Phys. Lett.*, 1990, 165, 107.
- 18 C. Song, K.-N. Hu, C.-G. Joo, T. M. Swager and R. G. Griffin, *J. Am. Chem. Soc.*, 2006, **128**, 11385.
- 19 J. Wang, L. Hou, W. R. Browne and B. L. Feringa, *J. Am. Chem. Soc.*, 2011, **133**, 8162.
- 20 D. P. Goldberg, J. Tesler, J. Krzystek, A. G. Montalban, L.-C. Brunel, A. G. M. Barrett and B. M. Hoffman, *J. Am. Chem. Soc.*, 1997, **119**, 8722.
- 21 M. M. Roessler and E. Salvadori, *Chem. Soc. Rev.*, 2018, **47**, 2534.

View Article Online
DOI: 10.1039/D0DT04149A

Dalton Transactions Accepted Manuscript

