Environmental control programs the emergence of distinct product ensembles from unconstrained chemical reaction networks.

Andrew J. Surman,^{#1} Marc Rodriguez Garcia,^{#1} Yousef M. Abul-Haija,¹ Geoffrey J. T. Cooper,¹ Piotr S. Gromski,¹ Rebecca Turk-MacLeod,¹ Margaret Mullin,² Cole Mathis,³ Sara I. Walker,³ and Leroy Cronin*¹

- 1) WestCHEM, School of Chemistry, University of Glasgow, Glasgow, UK.
- 2) School of Life Sciences, MVLS, University of Glasgow, Glasgow, UK.
- 3) Beyond Center for Fundamental Concepts in Science, Arizona State University, Tempe, AZ, USA.

*Corresponding author: Lee.Cronin@glasgow.ac.uk

Table of contents

1	Gei	neral experimental details	.3
1.1	Rea	gents	3
1.2	Ger	neral Conditions of RP-HPLC-MS Analysis	3
1.3	Am	ino Acids	4
1.4	Рер	tide synthesis	4
2	Env	vironment-Directed Amino Acid (AA) Condensation Experiments	5
2.1	Env	vironment-Directed AA Condensation Experiments: Synthesis	5
2.1	.1	Effect of Soluble Salts (G, A, H)	5
2.1	.2	Effect of Minerals (G, A, H)	6
2.1	.3	Effect of Mixing History (G, A, H)	7
2.1	.4	Effect of Mixing History (A, V, D)	8
2.2	Env	vironment-Directed AA Condensation Experiments: Product Analysis	9
2.2	.1	Untargeted LC-MS & fingerprinting analysis approach	9

4	Ref	erences60
3.4	.2	Inspection of Assembly/Aggregation using TEM57
3.4	.1	Recognition assay using ThT57
3.4	Env	vironment-Directed Complex Mixture Experiments: Functional Examination57
3.3	Env	vironment-Directed Complex Mixture Experiments: Product Analysis50
3.2	Env	vironment-Directed Complex Mixture Experiments: Synthesis
3.1	Spa	rk Discharge Mixture Preparation48
3	Env	vironment-Directed Complex Mixture Condensation Experiments48
2.3	.4	Observation of different properties of gels produced on addition of Ca^{2+} salts46
2.3	.3	Inspection of Assembly/Aggregation using TEM45
2.3	.2	Recognition assay using ThT44
2.3	.1	Reactivity testing using <i>p</i> NPA41
2.3	Env	vironment-Directed AA Condensation Experiments: Functional Examination41
2.2	.3	Sequence permutation distribution difference between populations
2.2	.2	Differing populations: Untargeted LC-MS analysis Results & Discussion11

1 General experimental details

1.1 Reagents

All solvents used in synthesis were HPLC grade or higher; all solvents used in LC-MS analyses were LC-MS grade (VWR). Glycine, L-Alanine, L-Aspartic Acid, L-Histidine, L-Valine, *p*-nitrophenyl acetate, sodium chloride, potassium chloride, lithium chloride, magnesium chloride, europium (II) chloride, fumed silica, montmorillonite, Goethite, and Thioflavin T were purchased from Sigma Aldrich. Alumina was purchased from Acros Organics. Copper (II) chloride was purchased from Lancaster/Alfa Aesar. Natrolite and quartz were obtained from Richard Tayler Minerals, Cobham, Surrey, England, and used (crushed in a Teflon ball mill) without further purification. Mica was obtained from Agar Scientific, and used (crushed in a Teflon ball mill) without further purification. "Nanovan" negative stain for TEM was purchased from Nanoprobes. Deuterium oxide was supplied by Goss Scientific. Spectra/Por® Float-A-Lyzer® G2 dialysis tubes were purchased from Spectrum Labs. Gas mixtures were supplied pre-mixed by the British Oxygen Company (BOC) and CK Special Gases Ltd.

1.2 General Conditions of RP-HPLC-MS Analysis

Reversed-phase LC were performed using a Dionex Ultimate 3000 system fitted with an Agilent Poroshell 120 EC-C18 (4.6 x 150 mm, 2.7 μ m) column. Samples were typically injected in 2-5 μ L aliquots and eluted with a linear gradient mixture of solvents A (water w/0.1% *v/v* formic acid) and B (acetonitrile w/0.1% *v/v* formic acid) over 26 mins as follows: 0 min – 0% B; 4 min – 0% B; 16 min – 70% B; 19 min – 100% B; 23 min – 0% B. The column oven was maintained at 30 °C. The LC system was coupled to a MS apparatus: a Bruker MaXis Impact instrument, calibrated for the 50 – 1200 Da range using sodium formate solution. The eluent stream was introduced directly into the source (no splitting) following the DAD detector, at a dry gas temperature of 200 °C. The ion polarity for all MS scans recorded was positive, with the voltage of the capillary tip set at 4800 V, end plate offset at –500 V, funnel 1 RF at 400 Vpp and funnel 2 RF at 400 Vpp, hexapole RF at 100 Vpp, ion energy 5.0 eV, collision energy at 5 eV, collision cell RF at 200 Vpp, transfer time at 100.0 μ s, and the pre-pulse storage time at 1.0 μ s. In any MS/MS experiments CID energies were optimised according to products (typically between 20 and 30 eV).

All data acquisition was controlled by the Compass software suite, with DCMS Link/Chromeleon XPress. More complex analyses were performed using bespoke scripts in the R Environment.¹ To facilitate this, data files were converted to the open .mzML format, using Proteowizard MSConvert.²

Samples were used directly from the synthesis procedures, as described. Where too much material was present (causing saturation of MS detector through excess signal), all samples in the series were diluted by a 1 in 10 dilution, to allow injection in the 2-5 μ l volume range while optimising MS signal. The instrument was calibrated before each set of analytical replicates (each of which was completed before progressing to the next analytical replicate).

1.3 Amino Acids

Where amino acids (AAs) are discussed, they are frequently identified using standard singleletter notation: A = alanine; D = aspartic acid; G = glycine; H = histidine; V = valine. All those incorporating stereocentres are the L- enantiomer.

1.4 Peptide synthesis

Peptide standards (for identification of different G_4A sequence permutations) were synthesised separately using a standard solid phase technique (Fmoc Ala and Gly Wang resin; coupling with DIC/HOBT and a TFA cleavage; Fmoc deprotection with 20% Piperidine/DMF) using a Biotage Initiator+ Alstra Petide Synthesiser. DIC and TFA were purchased from Sigma Aldrich, and protected amino acids and Wang resin were purchased from Activotec.

2 Environment-Directed Amino Acid (AA) Condensation Experiments

2.1 Environment-Directed AA Condensation Experiments: Synthesis

2.1.1 Effect of Soluble Salts (G, A, H)

In this set of experiments, one solution containing an equimolar amount of three different amino acids was reacted to different soluble salts under successive dehydration-hydration cycles.

1. A solution containing three different amino acids (G, A, H) was prepared to a final concentration of 0.033 M (each) and adjusted to pH=2.5 by adding HCl.

	Experiment Label										
Cycle	NaCl	KCl	LiCl	NH ₄ Cl	MgCl ₂	CuCl ₂	EuCl ₃				
1	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H				
2	H_2O	H_2O	H_2O	H_2O	H_2O	H_2O	H_2O				
3	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O				
4	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H				
5	H_2O	H_2O	H_2O	H ₂ O	H_2O	H_2O	H ₂ O				
6	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O				
7	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H	G+A+H				
8	H_2O	H_2O	H ₂ O	H ₂ O	H ₂ O	H_2O	H ₂ O				
9	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O	H ₂ O				

2. 7 different soluble salt solutions were prepared at a final concentration of 1 M.

3. 1ml of a 1 M soluble salt solution was added in cycle 1 to each individual experiment.

4. 3.5 ml of the amino acids solutions were added in cycles 1, 4 and 7.

5. 3.5 ml of HPLC water were added in cycles 2, 3, 5, 6, 8 and 9.

 Each dehydration-hydration cycle was performed on a multiwell hotplate at 130 °C for 12 h (a fixed arbitrary cycle time; all repeat reactions performed together to avoid error).

- 7. Once finished, all the samples were diluted by adding 6 ml of HPLC water.
- 500 μl were taken for LC-MS analysis. The remaining sample was dialysed with a G2 Float-a-lyser (500-1000 Da) cut-off (5 ml) for 20 h.
- 9. Once the dialysis was completed, the samples were left to freeze-dry for 48 h.
- 10. The solid product material was redissolved in 6 ml of water, filtered through 0.22 μm syringe filters, and stored at 4°C to be used without further treatment.

2.1.2 Effect of Minerals (G, A, H)

In this set of experiments, one solution containing an equimolar amount of three different amino acids was reacted to different minerals under successive dehydration-hydration cycles.

1. A solution containing three different amino acids (G, A, H) was prepared to a final concentration of 0.033 M (each) and adjusted to pH=2.5 by adding HCl.

	Experiment Label											
Cycle	Alumina	Montmorillonite	Mica	Goethite	Quartz	Natrolite	Silica					
1	G+A+H											
2	H_2O	H ₂ O	H_2O	H_2O	H_2O	H_2O	H ₂ O					
3	H ₂ O											
4	G+A+H											
5	H_2O	H_2O	H_2O	H_2O	H_2O	H_2O	H ₂ O					
6	H ₂ O											
7	G+A+H											
8	H_2O	H ₂ O	H ₂ O	H ₂ O	H_2O	H_2O	H ₂ O					
9	H ₂ O											

- 2. 0.2 g of a powdered mineral were added in cycle 1 to each individual experiment.
- 3. 3.5 ml of the amino acids solutions were added in cycles 1, 4 and 7.
- 4. 3.5 ml of HPLC water were added in cycles 2, 3, 5, 6, 8 and 9.
- 5. Each dehydration-hydration cycle was performed on a multiwell hotplate at 130 °C for 12 h (a fixed arbitrary cycle time; all repeat reactions performed together to avoid error).
- 6. Once finished, all the samples were diluted by adding 6 ml of HPLC water.
- 500 μl were taken for LC-MS analysis. The remaining sample was dialysed with a G2 Float-a-lyser (500-1000 Da) cut-off (5 ml) for 20 h.
- 8. Once the dialysis was completed, the samples were left to freeze-dry for 48 h.
- 9. The solid product material was redissolved in 6 ml of water, filtered through 0.22 μ m syringe filters, and stored at 4°C to be used without further treatment.

2.1.3 Effect of Mixing History (G, A, H)

In this set of experiments, three different amino acid solutions were added in a different order of addition under successive dehydration-hydration cycles.

1. Three individual solutions of amino acids (glycine, alanine, histidine) were prepared to a final concentration of 0.1 M and adjusted to pH=2.5 by adding HCl. A mixture of the three ("G+A+H") was prepared by mixing these solutions 1:1:1 (v/v).

	Experiment Label										
Cycle	G⇔A⇔H	G⇔H⇔A	A⇔G⇔H	A⇔H⇔G	H⇔G⇔A	H⇒A⇔G	G + A + H				
1	G	G	А	А	Н	Н	G+A+H				
2	H ₂ O										
3	H ₂ O	H ₂ O	H ₂ O	H_2O	H ₂ O	H ₂ O	H ₂ O				
4	А	Н	G	Н	G	А	G+A+H				
5	H_2O	H_2O	H_2O	H_2O	H_2O	H_2O	H ₂ O				
6	H ₂ O										
7	Н	А	Н	G	А	G	G+A+H				
8	H ₂ O	H ₂ O	H ₂ O	H_2O	H ₂ O	H ₂ O	H ₂ O				
9	H ₂ O										

3. 3.5 ml of a 0.1 M solution of each amino acid were added in cycles 1, 4 and 7.

- 4. In the experiment where a mixture of the three amino acid solutions was added together,1.16 ml of each amino acid solution was added.
- 5. 3.5 ml of HPLC water were added in cycles 2, 3, 5, 6, 8 and 9.
- 6. Each dehydration-hydration cycle was performed on a multiwell hotplate at 130 °C for 12 h (a fixed arbitrary cycle time; all repeat reactions performed together to avoid error).
- 7. Once finished, all the samples were diluted by adding 6 ml of HPLC water.
- 500 μl were taken for LC-MS analysis. The remaining sample was dialysed with a G2 Float-a-lyser (500-1000 Da) cut-off (5 ml) for 20 h.
- 9. Once the dialysis was completed, the samples were left to freeze-dry for 48 h.
- 10. The solid product material was redissolved in 6 ml of water, filtered through 0.22 μ m syringe filters, and stored at 4°C to be used without further treatment.

(n.b. A separate set of experiments were also performed in which, instead of dissolving material in a fixed volume of water, a 0.5 mg/ml solution was made up; this is referred to in Section 2.3.1 as 'Constant Concentration' solutions, "CC", rather than 'Constant Volume', "CV").

2.1.4 Effect of Mixing History (A, V, D)

In this set of experiments, three different amino acid solutions were added in a different order of addition under successive dehydration-hydration cycles.

 Three individual solutions of amino acids (alanine, valine, and aspartic acid) were prepared to a final concentration of 0.1 M. The pH of each solution was acidified to below pH 2.5 by adding 3 ml of 5 M H₃PO₄ followed by adjusting the pH to 2.5 by adding a minimum amount of 5M NaOH.

	Experiment Label									
Cycle	1	2	3	4	5	6	7			
1	A+V+D	D	D	V	V	А	А			
2		V	А	D	А	D	V			
3		А	V	А	D	V	D			

2. The order in which the different amino acid solutions were added was decided.

- 3. 3.5 ml of a 0.1 M solution of each amino acid were added in cycles 1, 2 and 3.
- 4. In the experiment where a mixture of the three amino acid solutions was added together,3.5 ml of each amino acid solution was added.
- Each single dehydration-hydration cycle was performed on a multiwell hotplate at 130 °C for 12 h (a fixed arbitrary cycle time; all repeat reactions performed together to avoid error).
- 6. Once finished, all the samples were dissolved in 1.5 ml of HPLC water (by vortex and sonication for 5 min).
- 7. 500 μl were taken for LC-MS analysis. The remaining sample was pH adjusted to pH=7.0-7.5 with 500 μl of 5 M NaOH then filtered using 0.22 μm nylon syringe filters and stored at 4°C for further analysis (TEM analysis, ThT assay and gel formation).

2.2 Environment-Directed AA Condensation Experiments: Product Analysis

2.2.1 Untargeted LC-MS & fingerprinting analysis approach

Each reaction (performed in triplicate) was analysed three times in LC-MS, giving a total of 9 repeats (3 experimental x 3 analytical repeats). A qualitative overview of product distribution *vs* LC-MS intensity was obtained using bespoke script, under the R environment,¹ with files input in the ".mzML" format, and the xcms library³ for data extraction and peak picking functions. The procedure was as follows (results in following sections):

- i. Input all data in groups (9 experiments, in 7 groups).
- ii. Independently 'pick peaks' (*i.e.* detect features in signal, identified by m/z and retention time (*rt*) coordinates and characterised intensity values for each sample). [*xcms* 'Centwave' algorithm; 25 ppm error; peak prefilter requiring 7 data points of intensity > 1000; S/N required \geq 3; scanrange excluding 'column wash' part of LC cycle to minimise 'noise' contributions]
- iii. 'Group' peaks/features observed in many experiments with the same m/z and rt. [using xcms grouping; bw = 15; mzwid = 0.005]
- 'Fill in' missing data. *i.e.* where particular peaks were absent in some samples, extract intensity values at same *rt* and *m/z* values from samples where they were present. [*xcms* 'fillPeaks' function]

This produced a complete table of coordinates (m/z, retention time) for hundreds/thousands of features, along with intensity data for each LC-MS analysis.

- v. Perform Principal Component Analysis (PCA) of intensity variation of these picked peaks between analyses. PCA implemented using the FactoMineR library (with scaling),¹¹ and the first three PCs plotted using the rgl library (version 0.96.0)¹² or Origin Pro 2016,¹³ with 'bubbles' plotted around each set of experiments (each environment) representing two standard deviations around their mean (using ellipse3d function from the rgl library).
- vi. Principal component discriminant function analysis (PC-DFA) was also performed (using the MASS library),¹⁴ using the first five principal components (these accounted for the overwhelming majority of variance in all cases, see Section 2.2.2). This facilitated sharper observation of the differences between product populations (plotting the first three DFs), but was qualitatively similar to the results of simple (unsupervised) PCA analysis.

Notes and variations on this process:

- No attempt at this stage was made to identify unknown products the intention of this analysis is to obtain an overview of product distribution, a 'fingerprint', since thorough quantification & identification of every species present is neither practical nor necessary.
- Given these aims, peak picking algorithm settings were deliberately not stringent, to include as many features as possible. We note that while some noise may have been included as a result, its effect is likely to have been negligible: this is demonstrated through the observation that qualitatively similar differentiation of populations is observed when product peaks are filtered to include only potential product peptide masses from the AAs used (see Figure S3) and of the systematic variation of several peaks (see Figures S5 to S7 for example EICs). Furthermore, LC-MS/MS analysis of some species to identify isomers (see Figure S27c for typical example) demonstrates that peptide products are present as expected.
- Isobaric species (those with the same mass) are not resolved in MS detection, and since chromatographic separation frequently did not completely resolve manifolds of isobaric species resulting from different sequence permutations (*e.g.* GGGAG, GGAGG, GAGGG), in many cases it is likely that several species may have been included in the same 'feature' manifested as broad manifolds of coeluting peaks. Since in many cases the shape (composition distribution) and size (amount of species present) of these features tends to vary in a robust (reproducible) manner, this is not problematic for the conclusions drawn.

2.2.2 Differing populations: Untargeted LC-MS analysis Results & Discussion

General observations

- In all experiments, analysis reveals that many product populations are clearly and consistently different as a result of the variation of reaction environment: this can be observed in PCA (Figure S3) and PC-DFA analysis (Figure S2), and in extracted ion chromatogram (see Figures S5 to S7 for selected examples, demonstrating reproducible differences) and peak intensity data (Figures S8 to S10).
- PCA yields qualitatively similar results to PC-DFA in demonstrating this, but with less sharp separation. That is, the populations which can be observed to be similar, and those which are clearly resolved, in plots of PCA (Figure S3) are generally those of which similar observations can be made in PC-DFA plots (Figure S2). That PC-DFA, a supervised technique, provides sharper resolution than PCA (an unsupervised technique) is unsurprising; the qualitative similarity reflects the robust and reproducible nature of the difference between populations.
- In all cases, plotting contributions (Figure S4) to the principal components demonstrates that population difference is not defined by a few 'key' features/species; instead, many provide similar (small) contributions.
- Since in most cases experimental repeats produced extremely similar results, in cases where results are not very similar (large 'bubbles') we suspect that this largely due to material loss during sample work-up (filtering; dialysis; filtering; dissolution), for example inconsistency in dialysis membranes. This is consistent with observations during work on these systems (e.g. LC-MS analysis of undialysed samples).
- When the feature list was 'filtered' to exclude all masses not corresponding to a plausible oligomer or the amino acids used (from a combinatorial list of possible peptide products from the AAs combined, as "Peptide mass product distributions", +/-0.01 Da), the resulting plots (Figure S3) are qualitatively broadly similar to those unbiased by product expectations (the same populations are resolved/unresolved), demonstrating the robustness of the approach and that differences result from 'real' condensation products, not analytical artefacts.

Experiments varying soluble salts present

(Section 2.1.1; MCl_x salts, where $M = Na^+$, K^+ , Li^+ , NH_4^+ , Mg^{2+} , Cu^{2+} , or Eu^{3+})

- Monovalent salts produced similar product distributions (those from Na⁺, K⁺, and NH₄⁺ unresolved in simple PCA; resolved by PC-DFA, but adjacent), except Li⁺.
- Li⁺ experiments produced a product distribution similar to that produced in the presence of Mg²⁺. Presence of Cu²⁺ or Eu³⁺ leads to distributions which are clearly distinct from other salts.

Experiments varying minerals present

(Section 2.1.2; Minerals: Alumina, Montmorillonite, Mica, Goethite, Quartz, Natrolite, Silica)

• Most of the reactions incorporating minerals yielded product ensemble distributions which were robustly distinguished in all analyses performed (supervised and unsupervised), except the alumina/quartz pair.

Experiments varying amino acid mixing history

(Section 2.1.3; Orders: all permutations of sequential addition of G, A, and H; Shorthand " $G \Rightarrow A \Rightarrow H$ " means G added first, followed by condensation cycles, followed by addition of A, followed by condensation cycles, followed by addition of H, followed by condensation cycles; Shorthand "G+H+A" means all amino acids added together

- Broadly, most of the analyses resolve the ensembles into three pairs (G⇔A⇔H & A⇔G⇔H; G⇔H⇔A & H⇔G⇔A; A⇔H⇔G & H⇔A⇔G), with the reaction in which all amino acids were added together clearly resolved from all. In PC-DFA some of these pairs are resolved (although clearly adjacent), but this separation is not robustly observed across all analyses.
- The reaction pattern is consistent with the trends observed in preliminary binary crossreactivity tests ("Intensity"=sum of MS intensity accounted-for by putative combinatorial products) where G/A hetero-oligomerisation clearly dominates. For example, products of G⇒A reactions are likely to resemble A⇒G if G/A heterooligomerisation rates are very much larger than either possible homo-oligomerisation. While our approach in this work has been non-deterministic, interested in observing difference, these observations point to the potential for deliberate 'programming', using modelling of rate measurements, however, as we observe that simple thermodynamic considerations are not adequate, this will require a more advanced approach.




Figure S1. Plot of data from preliminary cross-reactivity investigation for different G, A and H amino acid combinations. "Intensity" is the combined intensity corresponding to the masses of putative oligomeric products (trimer and larger) produced when reacted in simple binary mixtures in the same conditions as used in Section 2.




Figure S2. Plots of PC-DFA analysis (using first 5 PCs) of results from experiments changing (a) soluble salts present, (b) minerals present, and (c) amino acid mixing history; in each case 'spots' represent individual measurements & 'bubbles' represent two standard deviations around their mean. Analysis conducted in R, calculated and plotted using rgl library.




Figure S3. Plots of PCA analysis comparing the full product distributions (a-c) to 'peptide mass product distributions' (d-f), produced by filtering the feature list for putative peptide products. Data drawn from experiments changing (a & d) soluble salts present, (b & e) minerals present, and (c & f) amino acid mixing history; in each case 'spots' represent individual measurements & 'bubbles' represent two standard deviations around their mean. Analysis conducted in R, calculated and plotted using rgl library.




Figure S4. Plots showing the fraction of variance explained by the principal components in Figure S3 (a-c) and the distribution of contributions to the first two principal components: changing soluble salts present (a & b), changing minerals present (c & d), and amino acid mixing history (e & f).




Figure S5. Selected extracted ion chromatograms illustrating product distribution variance in products from experiments varying soluble salt present (ordered by ascending m/z). (a) m/z = 156.077; (b) m/z = 164.082; (c) m/z = 167.066; (d) m/z = 198.089; (e) m/z = 209.104; (f) m/z = 229.141; (g) m/z = 284.135; (h) m/z = 312.151; (i) m/z = 355.172; (j) m/z = 369.188 [lines = mean intensity from all 9 measurements (3 experimental replicates x 3 analytical replicates); shading around line represents one standard deviation around mean; intensities normalised relative to largest value in each plot]




Figure S6. Selected extracted ion chromatograms illustrating product distribution variance in products from experiments varying which mineral is present (ordered by ascending m/z). (a) m/z = 261.119; (b) m/z = 284.135; (c) m/z = 432.184; (d) m/z = 475.190; (e) m/z = 567.253; (f) m/z = 569.243; (g) m/z = 683.286; (h) m/z = 695.312 [lines = mean intensity from all 9 measurements (3 experimental replicates x 3 analytical replicates); shading around line represents one standard deviation around mean; intensities normalised relative to largest value in each plot]




Figure S7. Selected extracted ion chromatograms illustrating product distribution variance in products from experiments varying amino acid mixing history (ordered by ascending m/z). (a) m/z = 218.114; (b) m/z = 261.119; (c) m/z = 374.203; (d) m/z = 426.210; (e) m/z = 446.119; (f) m/z = 489.205; (g) m/z = 517.237; (h) m/z = 635.264; (i) m/z = 683.286 [lines = mean intensity from all 9 measurements (3 experimental replicates x 3 analytical replicates); shading around line represents one standard deviation around mean; intensities normalised relative to largest value in each plot]




Figure S8. The intensity of each of the features picked (on which PCA, etc was performed), expressed as a fractional difference from a mean for all the peaks within a set to visualise variation in data. [i.e. (Mean^{NaCl} – Mean^{AllSalt}) / Mean^{AllSalt}; feature m/z and rt coordinates unlabelled, ordered by ascending m/z from left to right; Note: Fractional intensities can obscure smaller variations in intensity values below mean]




Figure S9. The intensity of each of the features picked (on which PCA, etc was performed), expressed as a fractional difference from a mean for all the peaks within a set to visualise variation in data. [i.e. (Mean^{Mica} – Mean^{AllMin}) / Mean^{AllMin}; feature m/z and rt coordinates unlabelled, ordered by ascending m/z from left to right; Note: Fractional intensities can obscure smaller variations in intensity values below mean]




Figure S10. The intensity of each of the features picked (on which PCA, etc was performed), expressed as a fractional difference from a mean for all the peaks within a set to visualise variation in data. [i.e. $(Mean^{G+A+H} - Mean^{AllHist}) / Mean^{AllHist}$; feature m/z and rt coordinates unlabelled, ordered by ascending m/z from left to right; Note: Fractional intensities can obscure smaller variations in intensity values below mean]

m/z	RT	NaCl	KCI	LiCl	NH4CI	MgCl2	CuCl2	EuCl3
95.0614	1.47	81291	111777	27499	139067	49595	26716	39130
110.0721	2.27	438569	337910	3324	412821	1321	259433	5103
110.0723	1.48	4091141					289740	795382
111.0753 121.9674	1.48	249333 4440	203500	32437	440125	53526 221	16868	44732 29494
123.9658	1.38	722	351	269	439	247	336082	14816
131.9632	1.40	532	384	619	371	398	221911	8176
143.0819	2.91	18578	28539	1057	31230	114	229529	438
147.0730	1.41	597673	863446	19260	1148462	395	15177	1868
148.9546	1.37	476	492	572	533	313	394289	11515
150.9505	1.40	614	549	460	757	617	179467	5583
152.5805	1.41	372699	353268	4356	474570	82	1144	416
156.0769	2.49	107418	89710	735	85522	481	215217	563
156.0770	1.48	1667356	2654349	346691	3240184	544851	170734	462555
156.0788	1.77	1872180	1998466	159072	2253800	247273	245254	230923
157.0809	1.48	104102	158055	26000	183609	40486	14029	34027
167.0934	1.79	513367	133556	10181	372304	1136	167400	9762
175.5832	1.47	405052	1120653	7383	428945	168	5868	1046
182.9637	1.34	1604	334	457	398	76569	554	325
184.0718	1.97	9422	14020	2547	17693	174	173063	651
184.9620	1.34	877	240	1042	445	23125	1800	382
194.1185	7.80	12629 2783962	491586	14451 37181	12310	17354 8649	10645 1092272	13083 68293
195.0886 195.8985	1.80	1355	491586	962	2080047	309	10922/2	155830
195.8985	1.40	5648	1082	1209	745	812	2699	403399
195.9186	1.29	1231	273	1209	208	136	8076	155843
197.9988	1.41	5807	920	460	442	345	2491	452150
200.9744	1.28	309	319	651	252	95054	404	452150
201.9737	1.34	282	233	278	267	28106	349	201
202.9715	1.34	275	150	184	216	27424	243	133
204.0953	1.67	181351	466368	1949	114356	235	12478	638
209.1038	2.22	2519913	433848	43118	1778546	15176	1853248	120314
210.1061	2.22	244075	45924	5279	199250	1755	204394	14303
211.1027	1.79	244346	458588	2658	287230	407	17022	960
213.0987	1.60	606675	598152	68881	709448	7065	95742	12671
213.1463	14.53	28464	27997	28197	28487	28556	29122	30237
215.6014	1.33	578071	513889	1648	609688	188	603	511
218.9840	1.34	190	192	147	220	238911	1367	148
219.9865	1.34	646	275	142	481	67241	1092	158
220.9812	1.34	247	247	119	240	71035	319	177
227.0795	2.29	48567	63017	963	44381	90	417951	981
227.1156	1.67	397435	372189	55985	568910	4135	85754	9156
239.6130	1.81	454720	327138	1453	296123	109	7270	469
240.9169	1.28	17374	2312	933	1061	911	829	1505588
242.0015	1.34	174	155	81	166	39965	159	156
242.9182	1.28	19780	2721	1229	1136	960	982	457
244.1135	1.34	622346	594080	17413	624014 17866	302 16832	516	17355
246.2444 255.1087	13.85	17274	16973 16189	17413	17800	58	380745	233
258.9276	1.28	11363	1664	906	692	778	1095	1087083
260.8539	1.40	3484	682	420	370	465	14541	590439
260.9291	1.40	12554	1793	950	808	915	3225	1187401
270.1215	1.67	442474	310065	15465	421304	92	10403	2811
274.2761	15.33	32374	32178	31739	32512	31605	32463	33021
275.1235	1.50	1494631	799074	32794	2095374	4287	110282	65051
275.1283	1.51	977387	438472	37333	1187790	4418	80934	67382
276.9384	1.28	9044	1397	825	646	673	1559	818361
278.8650	1.40	2962	618	417	426	394	29052	503661
278.9399	1.29	9468	1325	918	658	732	7196	901200
281.9436	1.29	1377	358	592	273	464	2125	177332
283.9453	1.29	1548	320	401	285	357	1386	196250
284.1327	1.91	251775	237533	15810	341876	347	65069	4229
284.1378	1.91	442873	406552	13517	576827	160	47836	1555
293.1381	1.44	309414	310656	10110	462540	1762	5653	1289
294.9490	1.29	4964	577	556	375	586	1426	424602
296.9504	1.29	3215	458	277	301	208	245	321751
301.9558	1.29	1346	293	255	183	326	180	140891
311.1484	1.47	107596	26003	32899	112823	67145	601	48710
333.1184	1.48	8851	5855	2360	4019	29337	280	3963
339.7792	1.40	2112	3462	134	1318	36	241848	2374
341.7777	1.40	345	587	168	321	23	330433	3058
350.1550	1.41	733300	881125	6948	1267554	280	2781	863
350.1600	1.41	822775	917556	8072 8016	615381	299 409	3139 8036	915 2768
364.1755	1.50	66	445925	8016	615381	29005	63	2768
400.9430	1.34	352275	478800	1471	475317	118	1724	547
407.1804 421.1984	1.46 1.58	534376	518929	3345	612085	290	1724	1140
466.7091	1.38	2844	3006	66	1768	16	290545	324
			0000			10	20070	U.L.T

Figure S11. Table of selected features ordered by m/z from the experiments where the soluble salt present was varied. Features were selected from a full list based on absolute MS intensity (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	NaCl	KCI	LiCl	NH4Cl	MgCl2	CuCl2	EuCl3
242.9182	1.28	19780	2721	1229	1136	960	982	1705296
240.9169	1.28	17374	2312	933	1061	911	829	1505588
197.9189	1.28	5807	920	460	442	345	2491	452150
260.9291	1.28	12554	1793	950	808	915	3225	1187401
258.9276	1.28	11363	1664	906	692	778	1095	1087088
276.9384	1.28	9044	1397	825	646	673	1559	818361
195.9186	1.29	5648	1082	1209	745	812	2699	403399
278.9399	1.29	9468	1325	918	658	732	7196	901200
296.9504	1.29	3215	458	277	301	208	245	321751
294.9490	1.29	4964	577	556	375	586	1426	424602
281.9436	1.29	1377	358	592	273	464	2125	177332
301.9558	1.29	1346	293	255	183	326	180	140891
283.9453	1.29	1548	320	401	285	357	1386	196250
215.6014	1.33	578071	513889	1648	609688	188	603	511
200.9744	1.34	309	319	651	252	95054	404	454
201.9737	1.34	282	233	278	267	28106	349	201
220.9812	1.34	247	247	119	240	71035	319	177
202.9715	1.34	275	150	184	216	27424	243	133
184.9620	1.34	877	240	1042	445	23125	1800	382
			192					
218.9840	1.34	190	112101	147	220	238911	1367	148
244.1135	1.34	622346	594080	1654	624014	302	516	457
400.9430	1.34	66	55	77	77	29006	63	811
182.9637	1.34	1604	334	457	398	76569	554	325
219.9865	1.34	646	275	142	481	67241	1092	158
242.0015	1.34	174	155	81	166	39965	159	156
148.9546	1.37	476	492	572	533	313	394289	11515
121.9674	1.38	4440	3834	419	5065	221	678729	29494
123.9658	1.38	722	351	269	439	247	336082	14816
131.9632	1.40	532	384	619	371	398	221911	8176
260.8539	1.40	3484	682	420	370	465	14541	590439
195.8985	1.40	1355	437	962	383	309	10117	155830
150.9505	1.40	614	549	460	757	617	179467	5583
466.7091	1.40	2844	3006	66	1768	16	290545	324
468.7076	1.40	258	314	31	281	51	194447	240
339.7792	1.40	258	314	134	1318	36		240
Contraction of the second s		-					241848	-
278.8650	1.40	2962	618	417	426	394	29052	503661
341.7777	1.40	345	587	168	321	23	330433	3058
197.8988	1.41	1231	273	168	208	136	8076	155843
147.0730	1.41	597673	863446	19260	1148462	395	15177	1868
350.1600	1.41	822775	917556	8072	1280548	299	3139	915
152.5805	1.41	372699	353268	4356	474570	82	1144	416
350.1550	1.41	733300	881125	6948	1267554	280	2781	863
293.1381	1.44	309414	310656	10110	462540	1762	5653	1289
407.1804	1.46	352275	478800	1471	475317	118	1724	547
95.0614	1.47	81291	111777	27499	139067	49595	26716	39130
311.1484	1.47	107596	26003	32899	112823	67145	601	48710
175.5832	1.47	405052	1120653	7383	428945	168	5868	1046
111.0753	1.48	249333	203500	32437	440125	53526	16868	44732
110.0723	1.48	4091141	2842384	564957	8759391	954859	289740	795382
157.0809	1.48	104102	158055	26000	183609	40486	14029	34027
333.1184	1.48	8851	5855	2360	4019	29337	280	3963
156.0770	1.48	1667356	2654349	346691	3240184	544851	170734	462555
Contraction of the Property of the		1494631	799074	346691	2095374	4287	110282	65051
275.1235	1.50	(D)			10	4287		-
364.1755	1.50	321495	445925	8016	615381		8036	2768
275.1283	1.51	977387	438472	37333		4418	80934	67382
421.1984	2100	534376	518929	3345	612085	290	10250	1140
213.0987	1.60	606675	598152	68881	709448	7065	95742	12671
204.0953	1.67	181351	466368	1949	114356	235	12478	638
227.1156	1.67	397435	372189	55985	568910	4135	85754	9156
270.1215	1.67	442474	310065	15465	421304	92	10403	2811
156.0788	1.77	1872180	1998466	159072	2253800	247273	245254	230923
211.1027	1.79	244346	458588	2658	287230	407	17022	960
167.0934	1.79	513367	133556	10181	372304	1136	167400	9762
195.0886	1.80	2783962	491586	37181	2080047	8649	1092272	68293
239.6130	1.81	454720	327138	1453	296123	109	7270	469
284.1327	1.91	251775	237533	15810	341876	347	65069	4229
284.1378	1.91	442873	406552	13517	576827	160	47836	1555
184.0718	1.97	9422	14020	2547	17693	174	173063	651
209.1038	2.22	2519913	433848	43118	1778546	15176	1853248	120314
		an1	1				1000 C	
210.1061	2.22	244075	45924	5279	199250	1755	204394	14303
110.0721	2.27	438569	337910	3324	412821	1321	259433	5103
227.0795	2.29	48567	63017	963	44381	90	417951	981
156.0769	2.49	107418	89710	735	85522	481	215217	563
255.1087	2.54	12348	16189	190	17764	58	380745	233
143.0819	2.91	18578	28539	1057	31230	114	229529	438
194.1185	7.80	12629	10981	14451	12310	17354	10645	13083
246.2444	13.85	17274	16973	17413	17866	16832	17457	17355
	14.53	28464	27997	28197	28487	28556	29122	30237
213.1463			21331			20000	27122	

Figure S12. Table of selected features ordered by RT from the experiments where the soluble salt present was varied. Features were selected from a full list based on absolute MS intensity (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	NaCl	KCI	LiCI	NH4CI	MgCl2	CuCl2	EuCl3
204.0923	3.02	5181	6166	87	5424	52	8949	54
204.0953	1.67	181351	466368	1949	114356	235	12478	638
218.1094	1.93	122244	108328	1082	166526	221	12835	803
218.1134	2.62	14570	26775	428	29046	33	22173	117
261.1160	1.78	60499	86010	655	57240	160	4227	475
270.1215	1.67	442474	310065	15465	421304	92	10403	2811
275.1277	7.26	9387	8098	365	9174	259	2944	309
275.1283	1.51	977387	438472	37333	1187790	4418	80934	67382
275.1326	2.66	51860	55589	487	62228	299	10189	325
284.1327	1.91	251775	237533	15810	341876	347	65069	4229
284.1378	1.91	442873	406552	13517	576827	160	47835	1555
298.1533	2.25	44642	38966	486	66118	50	11997	443
327.1430	1.78	159347	142156	977	188869	96	5184	574
332.1482	1.36	133398	166117	944	152817	217	1056	259
341.1523	1.33	118881	142657	548	105014	152	7620	100
341.1565	2.17	160772	141577	779	160775	162	13724	395
341.1601	2.18	187583	154770	766	224560	160	39037	395
350.1550	1.41	733300	881125	6948	1267554	280	2781	863
350.1600	1.41	822775	917556	8072	1280548	299	3139	915
355.1677	1.59	31698	39450	166	14645	119	1720	103
355.1702	1.99	112959	74790	1216	76827	114	30833	638
355.1757	1.92	149316	124614	1290	166874	117	26526	675
364.1755	1.50	321495	445925	8016	615381	409	8036	2768
369.1904	2.78	14769	14560	182	18881	83	10212	184
384.1650	1.64	223278	199601	722	234681	68	2175	790
398.1741	1.87	112325	103148	554	193086	118	4973	346
398.1789	1.89	194539	179934	1036	199310	225	11920	824
407.1804	1.46	352275	478800	1471	475317	118	1724	547
417.2034	2.43	14933	2607	67	11394	35	9800	158
421.1984	1.58	534376	518929	3345	612085	290	10250	1140
430.1945	1.37	136750	114699	441	147853	192	149	117
430.1989	1.30	135843	127486	409	147897	178	124	116
435.2125	1.70	103067	105607	650	110105	63	9635	482
487.2180	1.33	272083	237889	345	278866	96	160	278
567.2564	1.24	12303	9566	38	15493	36	26	598
638.2966	1.26	11334	15619	48	14555	66	31	1314

Figure S13. Table of selected features ordered by m/z from the experiments where the soluble salt present was varied. Features were selected from a full list based on absolute MS intensity, filtered to include masses consistent with peptides, (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	NaCl	KCI	LiCI	NH4CI	MgCl2	CuCl2	EuCl3
567.2564	1.24	12303	9566	38	15493	36	26	598
638.2966	1.26	11334	15619	48	14555	66	31	1314
430.1989	1.30	135843	127486	409	147897	178	124	116
341.1523	1.33	118881	142657	548	105014	152	7620	100
487.2180	1.33	272083	237889	345	278866	96	160	278
332.1482	1.36	133398	166117	944	152817	217	1056	259
430.1945	1.37	136750	114699	441	147853	192	149	117
350.1600	1.41	822775	117556	8072	280548	299	3139	915
350.1550	1.41	733300	881125	6948	1267554	280	2781	863
407.1804	1.46	352275	478800	1471	475317	118	1724	547
364.1755	1.50	321495	445925	8016	615381	409	8036	2768
275.1283	1.51	977387	438472	37333	1187790	4418	80934	67382
421.1984	1.58	534376	518929	3345	612085	290	10250	1140
355.1677	1.59	31698	39450	166	14645	119	1720	103
384.1650	1.64	223278	199601	722	234681	68	2175	790
204.0953	1.67	181351	466368	1949	114356	235	12478	638
270.1215	1.67	442474	310065	15465	421304	92	10403	2811
435.2125	1.70	103067	105607	650	110105	63	9635	482
327.1430	1.78	159347	142156	977	188869	96	5184	574
261.1160	1.78	60499	86010	655	57240	160	4227	475
398.1741	1.87	112325	103148	554	193086	118	4973	346
398.1789	1.89	194539	179934	1036	199310	225	11920	824
284.1327	1.91	251775	237533	15810	341876	347	65069	4229
284.1378	1.91	442873	406552	13517	576827	160	47835	1555
355.1757	1.92	149316	124614	1290	166874	117	26526	675
218.1094	1.93	122244	108328	1082	166526	221	12835	803
355.1702	1.99	112959	74790	1216	76827	114	30833	638
341.1565	2.17	160772	141577	779	160775	162	13724	395
341.1601	2.18	187583	154770	766	224560	160	39037	395
298.1533	2.25	44642	38966	486	66118	50	11997	443
417.2034	2.43	14933	2607	67	11394	35	9800	158
218.1134	2.62	14570	26775	428	29046	33	22173	117
275.1326	2.66	51860	55589	487	62228	299	10189	325
369.1904	2.78	14769	14560	182	18881	83	10212	184
204.0923	3.02	5181	6166	87	5424	52	8949	54
275.1277	7.26	9387	8098	365	9174	259	2944	309

Figure S14. Table of selected features ordered by RT from the experiments where the soluble salt present was varied. Features were selected from a full list based on absolute MS intensity, filtered to include masses consistent with peptides, (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

1100713 7.41 34109 6121.44 6427 7289895 43339 1172 92020 110072 1.48 20309 7/47 10007 24107 9709 10007 125,077 1.40 3574 4461 1009 701 12127 0404 9709 1007 125,077 1.40 3574 4461 1009 70 12129 0404 9709 1209 0409 1007 1219 0404 1209 0405 1007 1219 0409 1007 1219 0409 1007 1009 1007 1009 <th>m/z</th> <th>RT</th> <th>Alumina</th> <th>Montmoril.</th> <th>Mica</th> <th>Goethite</th> <th>Quartz</th> <th>Natrolite</th> <th>Silica</th>	m/z	RT	Alumina	Montmoril.	Mica	Goethite	Quartz	Natrolite	Silica
1110715 18.1 10.2586 0.7917 0.46600 10.000	95.0607	1.47	145157	289066	138974	342833	146885	38298	234199
1110742 1.4.8 22380 7787 1.9097 1.9087 1.9107 1.9107 1.9107 1.9007 1.9107 <th>110.0713</th> <th>7.41</th> <th>364164</th> <th>351148</th> <th>84117</th> <th>568845</th> <th>435159</th> <th>11725</th> <th>958289</th>	110.0713	7.41	364164	351148	84117	568845	435159	11725	958289
144.7716 1.0 19:00 3100 19:00 3104 19:01 19:02 99:00 10:00 152.5797 1.0 133 137 44 69:7 121.5 12277 14:00:09 1228 12277 14:00:09 1228 12297 14:00:09 126 12277 14:00:09 126 12277 14:00:09 126 125 126 126 12277 14:00:09 126 126 126 126 126 126 126 126 126 126 126 126 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 126 126 127 128 128 127 126 12	110.0715	1.81	1035866	687981	446650	1142986	2783821	131242	1759590
152.5797 1.0 1001	111.0742	1.48	225188	77487	150542	79025	241627	39939	165199
154.0073 1.3 17 44 695 210 1227 144000 0527 155.0765 1.59 1000 7981 10101 10001 6600 155.0765 1.59 1000 7981 10101 10001 10001 100000 100000 100000 100000 100000 100000 100000 100000 100000 1000000 10000	147.0716	1.40	740358	51620	398704	10448	749182	9076	350371
156.0764 7.40 1009 99.07 97.2 12.120 14.4071 1009 99.07 156.0767 3.77 22.04 27.05 27.04 <td< th=""><th>152.5797</th><th>1.40</th><th>357541</th><th>16051</th><th>134391</th><th>3551</th><th>341267</th><th>6916</th><th>128251</th></td<>	152.5797	1.40	357541	16051	134391	3551	341267	6916	128251
156.0765 15.9 20.84 <	154.9073	1.33	337	441	6695	210	12237	140056	2254
155.0797 3.77 2.50 9.70 1.61 9.90 1.64 9.90 1.62 157.0793 1.64 1.7960 1.386 2.2014 2.9044 3.930 3.930 1.933 168.0766 1.44 1.7960 1.386 2.2014 2.9044 3.930 3.940 1.237 175.0264 1.43 4.930 2.401 2.40	156.0764	7.40	185999	99617	67575	121150	248873	12003	664988
157.0793 1.6.5 15.04 4.184 9031 10092 9090 1122 166.0057 1.64 1976 2.454 3.107 9141 3.108 1.101 175.074 1.63 3.097 2.4573 3.097 9141 3.108 3.486 1.178 175.074 1.63 3.097 2.0177 910 4.0077 7.79 2.127 178.0764 5.33 3.78 3.441 4.13 3.147 2.779 2.0177 910 4.0077 7.79 2.127 180.0764 6.38 4.144 1.179 3.13 2.779 2.0177 910 4.0077 7.79 2.127 180.0765 2.955 1.170 7.120 7.77 9.177 <t< th=""><th>156.0765</th><th>1.59</th><th>2024089</th><th>724691</th><th>1831942</th><th>1023374</th><th>2309911</th><th>41<mark>9797</mark></th><th>1880253</th></t<>	156.0765	1.59	2024089	724691	1831942	1023374	2309911	41 <mark>9797</mark>	1880253
isf. 0025 1.6.4 1.986 1.2014 9.994 1.948 9.702 9.914 156.0768 1.6.3 6.037 3.9565 3.007 1.9319 2.9444 1.177 175.0274 1.6.3 6.037 2.0271 2.9221 4.114 4.781 2.9041 3.2441 3.2441 4.781 2.9071 3.2441 4.781 2.9091 3.331 2.272 3.91 3.918 3.917 4.9181 3.918 3.918 4.918 3.917 3.918 4.918 3.917 3.918 4.918 3.917 3.918 4.918 3.918 3.918 4.918 3.918 3.918 4.918 3.918 3.917 3.918 4.918 3.918 3.917 3.918 4.918 3.918 3.917 3.918 3.918 3.917 3.918 3.918 3.918 3.917 3.918 3.918 3.917 3.918 3.917 3.918 3.917 3.918 3.918 3.917 3.917 3.9177 3.9188 3.917<	156.0767	3.77	21543	62783	7596	574 <mark>864</mark>	22279	1512	69405
iss.orgs 1.4.4 1997 2.9466 3197 2.9437 9130 2.9466 92.771 175.0974 1.6.3 60.00 2.9271 2.9217 549.0 4.071. 2.949.1 2.949.1 175.0974 1.33 3.922 2.93 2.917.7 549.0 4.957.4 2.929.1 2.917.1 549.0 4.957.4 2.917.1 549.0 4.957.4 2.917.1 549.0 4.957.4 2.917.1 549.0 549.1 3.917.1 549.1 3.917.1 3.9	157.0793	1.65	155643	64398	114414	66553	150392	29600	122255
17.5.0974 16.3 60.09 2/771 2/221 4114 4/74 2/001 2/212 175.5824 1.68 5/60% 2/212 2/017	167.0925	1.64	173668	138856	220184	59014	198455	7332	91836
175.8242 1.48 54005 2077 2017 5400 40457 7200 2012 180.0764 5.35 903 2441 441 141 1424 2014 540 <th>168.0768</th> <th>1.44</th> <th>19697</th> <th>345645</th> <th>31007</th> <th>453617</th> <th>19130</th> <th>24846</th> <th>117762</th>	168.0768	1.44	19697	345645	31007	453617	19130	24846	117762
17.9.9021 1.3.3 2.22 49 33.8 119 6.49 7.90% 139 180.0764 6.38 43.14 4423.2 377 134.18 2424 46.3 4431 180.0764 6.38 43.14 4423.2 377 134.18 2424 46.3 4431 180.0765 2.05 1707 772.2 378 771.3 195.7 208.0 4490 181.0021 1.33 2.05 4.06 226.7 210.56 244.7 227.0 4.960 181.0021 1.33 3.62 4.07 20.22 210.56 34.04 172.22 210.44 182.0021 1.33 3.962 2.84.12 211.8 3.95.05 36.00 4.95.0 195.0661 1.80 2.87.7 1.96.15 2.96.0 9.96.94 1.99.0 4.96.0 195.0665 2.90 2.95.05 9.96.55 9.96.96 1.99.0 37.0 9.92.0 195.0666 2.79	175.0974	1.63	81035	247721	242281	48134	42781	20891	244586
180.0763 5.55 903 2041 941 91724 2122 944 9174 180.0765 2.55 19729 7120 9126 7131 1453 9127 180.0765 2.55 19729 7120 9126 7131 1453 9127 180.0075 1.33 925 60 924 127437 92762 1486 181.0072 1.33 962 417 414 2276 117437 12783 1284 182.0917 1.49 1000 1000 1100 7778 1013 1124 1272 11391 1244 184.0717 1.98 2020 24212 12406 1100 7785 1033 1344 1272 1333 1455 1033 1344 1214 1000 1343 144 1114 1000 1414 401 1215 10356 1036 10493 11980 1414 401 1199 120105 12210 121010	175.5824	1.48	546595	27370	201717	5460	404957	7829	213271
180.0764 6.38 4.14 4.422 177 1.3418 294 4.60 4.701 180.0765 2.55 1727 1.23 226 777 9170	179.9021	1.33	222	349	3436	1159	6429	76960	1260
180.00765 2.95 197/29 2120 2128 198/20 7131 14.53 2131 180.0015 1.33 2.95 4.00 7377 91702 19729 2400 181.0025 1.33 9.62 877 2177 19720 19729 2400 182.0921 1.43 1.642 14000 21222 21086 24171 1722 11729 121729 121721 121729 12181 184.0717 1.98 1262 2141 2118 3553 1841 1202 1131 1204 1205 3847 2211 1211 1204 1206 1208 3840 422 580 196.0661 1.80 287.75 161615 2016 1208 4804 11583 4233 422 580 200.066 2.29 1205.05 1211.010 120 4208 400 980 200.00 2002.0 2002.0 2002.0 2002.0 2002.0 2002.0	180.0763	5.35	3083	26431	941	343724	2102	504	19676
180.0015 1.33 235 4.00 1994 256 177 1702 1400 181.007 2.08 9071 1550.017 1530.017 1530.017 1530.017 1530.017 1530.017 1530.017 1537.02 2133 1607 6178 2076 11377 13732 2134 182.9027 1.33 166 2777 4666 1180 7787 160 103 1033 1031 1133 184.0717 1.98 2502 21412 2118 1550.53 12020 4001 0103 1134 196.0669 2.29 2194.64 940.55 1206.00 12020 4023 4020 402 402 4020 <	180.0764	6.38	4334	44252	1767	424188	2994	463	47418
180.0015 1.33 235 4.00 1994 256 177 1702 1400 181.007 2.08 9071 1550.017 1530.017 1530.017 1530.017 1530.017 1530.017 1530.017 1537.02 2133 1607 6178 2076 11377 13732 2134 182.9027 1.33 166 2777 4666 1180 7787 160 103 1033 1031 1133 184.0717 1.98 2502 21412 2118 1550.53 12020 4001 0103 1134 196.0669 2.29 2194.64 940.55 1206.00 12020 4023 4020 402 402 4020 <			10720	71250	3826	660407	7113	1453	38173
IB:1078 2.08 9071 13544 22347 3853 11747 2909 44900 IB:10025 1.33 1.49 1.304 1.0055 2122 20156 24374 1.722 2754 IB:0017 1.89 1.33 1.405 277 4068 1100 77815 6001 1101 IB:0017 1.88 0.771 1.88 0.771 1.88 0.771 1.88 0.771 1.88 0.771 1.88 0.771 1.80 2.785 0.8058 3.300 6.911 0.775 6.8013 0.90589 1.9559 0.902			255	430	3994		7377	91702	1405
181.9025 1.33 90.0 87 01.74 2.09 1177 1177 11792 2.141 182.9027 1.33 1405 2779 4066 1180 7783 1072 1131 11722 2184 184.0717 1.98 3262 21413 2118 35253 34872 34873 1801 11341 195.0876 1.80 2075 14815 21389 3486 190500 4000 602 9000 4030 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 60314 11390 11390 11390 11390 11390 11390 11390 11390 11390 11390 11390 11390 11390 11390 11393 1139				152646	225367			2490	449602
182.0017 1.49 1384 19930 2122 20196 2424 1722 7793 1925 1313 188.077 1.98 2.9022 21422 2118 35253 38407 20513 11541 195.0876 1.80 2.827 2148 35253 38407 20513 11541 195.0876 1.80 2.877 13051 4644 141804 4203 62 42743 198.0868 3.30 6111 11501 4644 141804 4203 62 5000 2790 42764 19700 1780 19707 1780 18707 1780 18707 1780 18707 1780 18707 1870 18707 1870 18707 18708 18707 18707 18707 18707 18707 18707 18708 18707 18708 18707 18708 18707 18708 18707 18708 18707 18708 18707 18708 18707 18708 18707 <th></th> <th></th> <th>land a second se</th> <th>887</th> <th>6174</th> <th>2370</th> <th>11727</th> <th>137192</th> <th>2148</th>			land a second se	887	6174	2370	11727	137192	2148
182.0022 1.33 140 277 4994 110 7799 100 1133 188.00717 1.98 32602 21429 2118 195.087 1.80 22020 1140 12020 1140 105669 110022 14010 10202 14010 10202 14010 10202 14010 10202 10410 10202 10410 10202 10410 10202 10410 10202 10410 10202 10410 10202 10410 10202 <th></th> <th></th> <th></th> <th>-</th> <th></th> <th></th> <th></th> <th>1</th> <th>27646</th>				-				1	27646
184.0717 1.98 33200 214.27 211.88 55253 3347 9031 11441 195.0876 1.80 20257 1.8015 21189 31968 1.8250 34210 7283 196.0901 1.80 20275 1.8015 21189 31968 1.6250 560 42343 198.0868 3.30 6111 11501 4644 1.41804 4233 612 560 42343 100.066 2.29 220168 64649 1.9431 64649 1.990 3.900 2.900 3.900 3.900 3.970 5207 211.0666 4.66 1.99 2.868 541 1.9481 1.964 4.90 4.90 211.001 1.72 2.9807 1.8722 1.7880 1.9766 3.9968 2.937 1.9354 1.9368 4.91 1.9356 4.91 1.9356 3.9368 4.91 1.9357 1.1369 3.937 1.9364 1.935 3.9356 3.9356 3.9356								-	13335
195.0876 1.80							33487	30513	119416
196.0901 1.80 227.75 1.81.31 211.349 1.31.461 1.95.55 50.9 4.94.34 198.0868 3.30 4111 1.15.61 644.4 1.14.84 4.20.31 51.2 50.30 209.0669 2.29 1.21.56.8 644.51 1.14.84 4.60.1 1.19.90 2.27.5 1.60.66 210.0154 2.23 1.44.27 1.14.87 1.34.83 660.14 1.19.90 2.27.5 1.60.66 210.0154 2.23 1.44.27 1.14.87 1.27.2 2.27.66 4.66.0 1.99.902 3.17.7 1.27.2 1.19.66 4.66.1 1.99.97 1.77.8 3.11.70 6.32.2 1.66.83 5.18.8 1.12.02 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.80 2.33.7 1.13.83 2.33.7 1.13.83			3560203	2007331	2897008	390669	2019202	34010	5572833
198.0868 3.30 6111 11301 6444 14104 203 612 500 209.0669 2.29 129153 101683 11943 66164 11970 8.780 200.1030 2.22 11843 11843 11963 11963 6050 12075 <th></th> <th></th> <th>287575</th> <th>163615</th> <th>231349</th> <th>31968</th> <th>163565</th> <th>3560</th> <th>457434</th>			287575	163615	231349	31968	163565	3560	457434
209.0669 2.29 2.21 2.21 2.23 2.23 2.23 2.24 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.243 2.245 2.245 2.245 2.245 2.245 2.245 2.245 2.245 2.245 2.2568 4.660 2.9597 1.772 2.7598 4.660 2.0597 1.374 2.2571 1.1878 2.1064 2.203 2.133 2.1350 2.2571 1.1878 2.205 2.1318 2.203 2.233 1.1884 2.27.0769 2.30 2.222 4.9363 1.5567 5.3544 2.2397 1.1854 2.24.0928 2.05 2.2117 1.222 1.514 4.88 2.8944 1.99 1.9252 2.44.0928 2.05 2.1817 1.225 1.534 1.077 1.191 1.182 2.44.0928 2.055									56031
209.1030 2.22 2.0302 2.0433 2.0303 3.0000 2.0399 2.2793 2.0000 210.1054 2.23 346576 111170 2.7068 4.600 1.90902 3.370 9.5277 210.0666 4.66 1.999 2.8688 9.41 9.4218 1.104 4.09 9.805 211.0606 1.72 2.9607 1.472 7.780 1.8768 9.3956. 4.001 1.0254 213.0977 1.78 31076 6.522 1.6688 2.0766 5.337 9.13340 1.21052 1.13340 9.3855 227.1133 1.67 2.79633 87500 1.7567 5.3876 1.2534 1.283 9.4833 239.6115 1.77 3.3992 1.1222 1.0141 2.44 9.9842 3.997 1.919 244.0928 2.05 2.817 1.2525 1.859 1.4107 2.7714 4.423 9.9963 246.2419 1.3.4 9.843 9.122 5.19877					1016808		689164	11990	878018
210.1054 2.23 948376 181170 275088 40840 195902 3370 53270 211.0610 1.72 219607 14722 7780 31966 4001 409 3050 213.0977 1.78 111070 82522 16633 5106 11202 11340 5517 215.0010 1.33 304652 5522 92751 1542 33554 2333 1384 227.0769 2.30 2522 4330 10608 20076 1534 1533 1384 227.0769 2.30 2522 101541 20076 15344 1533 1384 240.028 2.05 21817 13535 1767 33834 2597 1354 246.2419 13.84 34852 3402 3355 410070 1199 1352 246.8617 1.36 299 672 419 32553 204042 1417 1276 264.8617 1.36 299			3893142	2101418		380200	1969699	28795	6189056
211.0686 4.66 1599 28368 941 442118 1604 409 9850 211.1010 1.72 219607 1.472 7780 11788 33966 4001 1054 213.0977 1.78 311076 8222 7780 11788 33966 4001 1054 215.6010 1.33 30452 5532 92731 1542 326540 2333 1384 227.1133 1.67 177833 8759 15557 53876 33834 2537 18558 239.6115 1.77 33982 11252 10141 2438 24642 1999 13958 241.0928 2.05 2.117 13253 13859 141070 2104 4423 9054 246.2419 13.84 3452 34129 33525 3352 41070 1199 1362 266.8617 1.36 399 677 372 37534 3093 135 176 266.				-		1			552761
211.1010 1.72 219677 14722 77880 13788 33956 4001 10254 213.0077 1.78 311076 85522 16683 5108 121052 113800 15175 215.001 1.33 90452 5522 2731 1542 316564 2333 1384 227.0769 2.30 29228 49306 19765 55876 55876 55874 1593 26237 1853 239.6115 1.77 33992 11252 11514 248 25835 1070 1199 1352 244.0928 2.05 21877 13353 1859 17407 2704 4429 9952 246.817 1.36 399 872 419 3253 2090 352 244 246.8617 1.36 499 872 419 3253 2090 352 244 269.1241 1.66 9054 50977 37514 1603 155 176									
213.0977 1.78 311076 65252 166383 56108 212652 113360 35175 215.6010 1.33 304632 5532 92751 1542 336540 2333 1384 227.0769 2.300 29228 49306 17567 53876 358354 25377 1835 227.1133 1.67 379633 67850 17567 53876 358354 25377 1835 240.028 2.05 21817 13325 1859 314107 27044 4429 1999 1385 244.6113 1.36 6170 104192 2535 410070 1199 13662 246.6217 1.36 999 872 419 32533 2090 532 244 246.8731 1.36 424 667 372 37534 1603 155 176 267.1082 1.53 2662 1667 372 37534 1633 157 176 269			10					4001	102547
215.6010 1.33 304632 552 92731 1542 336540 2333 13844 227.0769 2.30 92922 49906 16008 20076 15244 1581 24433 227.1133 1.67 336952 11252 101541 2438 224942 1900 13536 241.0928 2.05 2117 11252 101541 2438 24942 1900 13537 244.018 1.34 381586 6107 104192 3355 410070 1199 13627 246.8617 1.36 399 872 419 32533 2090 552 244 246.8617 1.36 499 872 419 32534 1003 155 176 267.1082 1.53 2662 1599 1881 1163 1603 155 176 269.1241 3.76 9455 7860 2128 248473 3111 607 60111 269.1241 5.35 5368 36910 1768 24259 4103 507 3						in the second se			351759
227.0769 2.30 2923 4996 16968 200786 15244 1583 2432 227.1133 1.67 379633 17750 17857 53876 338344 23937 1854 239.6115 1.77 338982 11322 101511 2448 24442 1999 13858 241.0928 2.05 21817 132535 1859 174107 27104 4429 90453 244.1118 1.34 38156 6.170 104192 3322 3324 33949 34868 3222 246.8617 1.36 399 872 419 32233 2090 152 244 248.1140 1.66 90834 509572 51967 287272 72684 14117 12769 264.8731 1.36 424 667 372 37534 1603 155 176 267.1082 1.53 2662 168 181 1145 3661 9739 2434						-			138847
227.1133 1.67 379633 87850 175657 53876 358354 25397 18354 239.6115 1.77 338982 1122 101541 248 289492 1099 13958 244.1118 1.34 38158 6170 104192 3535 410070 1199 1362 246.2419 13.84 34852 34129 33825 33828 33949 34668 3222 246.8617 1.36 399 672 419 3252 2044 117 12769 264.8731 1.36 424 667 372 37534 1003 155 176 267.1082 1.53 2662 1649 181 114 1003 155 176 269.1241 3.76 9435 76840 2128 53933 3111 67 633 270.1189 1.71 479694 5720 25165 1269 5416 726 2558 275.1253									248325
239.6115 1.77 338982 11252 101541 2438 284942 1999 13958 241.0928 2.05 21817 11253 18589 174107 27104 4429 9965 244.1118 1.34 381586 6170 104192 3535 410070 1199 1362 246.2419 1.3.84 34452 34129 33825 33829 34668 3222 246.8617 1.36 399 872 419 32533 2090 352 2444 248.1140 1.66 90844 509572 51967 27227 72844 14117 12769 267.1082 1.53 2662 1649 1881 1145 3661 97539 2443 269.1241 5.35 536 36910 1708 49729 416 775 3384 269.1241 5.35 536 36910 1708 49729 5416 7257 3384 2092 25183 1285 3424 2025 274276 1395 36475 23944 13956 <th></th> <th></th> <th>379633</th> <th>in the second se</th> <th></th> <th>10</th> <th>358354</th> <th>25397</th> <th>183543</th>			379633	in the second se		10	358354	25397	183543
241.0928 2.05 21817 112535 1889 174107 27104 4429 90453 244.1118 1.34 381566 6170 104192 3352 410070 1199 13622 246.6419 13.84 34852 94129 33625 3353 2000 352 244 246.8617 1.36 399 872 419 32553 2000 352 244 248.1140 1.66 9084 509572 51967 287227 7284 14117 12769 266.1082 1.53 2662 1649 181 1145 361 9739 2443 269.1241 3.76 9435 7680 2128 9333 3111 607 60111 269.1241 5.35 5368 36910 1788 37599 4103 507 3283 270.1189 1.71 47694 57820 21685 12569 5472 3065 20645 274.2736 <th></th> <th></th> <th>338982</th> <th>11252</th> <th>101541</th> <th>2438</th> <th>284942</th> <th>1909</th> <th>139587</th>			338982	11252	101541	2438	284942	1909	139587
244.118 1.34 381386 6170 104192 3335 410070 1199 13682 246.2419 13.84 34852 3129 33625 3828 33949 34688 3222 246.8617 1.36 399 872 419 32553 2090 352 244 248.1140 1.66 90844 50572 51997 28727 72884 14117 12769 264.8731 1.36 4.24 667 372 27514 1603 155 176 267.1082 1.53 2662 1649 1881 1145 3661 97539 2443 269.1236 6.38 4850 45283 2135 53933 3111 607 60111 269.1241 5.35 5368 36910 1708 47269 4103 507 3283 270.1189 1.71 470984 57820 251635 12659 584752 30845 20025 274.2736<			21817	132535	18589	174107	27104	4429	90458
246.2419 13.84 34852 34129 33625 33828 33949 34668 32922 246.6617 1.36 399 872 419 32553 2000 352 244 248.1140 1.66 90834 509572 51967 287227 72684 14117 12769 264.8731 1.36 424 667 372 37514 1603 155 176 267.1082 1.53 2662 1649 1881 1145 3661 97539 243 269.1236 6.38 4850 45283 2195 53933 3111 607 6011 269.1241 5.35 5368 36910 1708 47269 4103 507 3283 270.1189 1.71 47694 57820 251635 12659 584752 30645 2040 282.8282 1.50 1490287 320607 448873 36275 133234 13956 12640 28.					104192	3535	410070	1199	136826
246.8617 1.36 399 872 419 23253 2090 352 244 248.1140 1.66 90834 509572 519967 287227 72684 14117 1269 264.8731 1.36 424 667 372 375514 1603 155 176 267.1082 1.53 2662 1649 1881 1145 3661 97539 244 269.1241 3.76 9435 76840 2128 709640 5416 726 3959 269.1241 5.35 5368 36910 1708 497269 4103 507 2283 270.1189 1.71 47694 57820 221635 12659 54752 30845 20805 274.2736 15.32 73172 7342 73607 132344 13956 126400 288.828 1.35 438 589 344 370686 1403 148 13956 284.1340 1.90					i i			0	32923
248.1140 1.66 90834 509572 519967 287227 72684 14117 12769 264.8731 1.36 424 667 372 375514 1003 155 176 267.1082 1.53 2662 1649 1881 1145 3661 97539 2433 269.1236 6.38 4850 45283 2128 109830 5416 726 6011 267.1082 109830 5416 726 6013 2019330 5416 726 6013 507 3283 269.1241 5.35 5368 36910 1708 497269 4103 507 3283 270.1189 1.71 476984 57820 251635 12659 584752 30845 20825 274.2736 15.32 73172 7342 7360 75945 74789 75705 7377 275.1253 1.50 190287 320607 24873 36275 132324 13956 12640			399		419	432553	2090	352	244
264.8731 1.36 424 667 372 375514 1603 155 176 267.1082 1.53 2662 1649 1881 1145 3661 97339 2443 269.1236 6.38 4850 45283 2155 53933 3111 607 60111 269.1241 3.76 9435 76840 1218 09840 5416 726 3958 269.1241 3.76 9435 76840 1218 09840 5416 726 3958 270.1189 1.71 476984 57820 251635 12659 564752 30845 20825 274.2736 15.32 73172 77482 73860 75945 74789 75705 73737 275.1253 1.50 1490287 320607 48873 36275 132334 13956 12640 284.8340 1.90 336839 39968 157867 13967 320152 8842 12785			90834	509572	519967	287227	72684	14117	1276975
267.1082 1.53 2662 1649 1881 1145 3661 97539 2443 269.1236 6.38 4450 45283 2195 53933 3111 607 6011 269.1241 3.76 9435 76840 2128 198840 5416 726 3385 269.1241 5.35 5368 36910 1708 417269 4103 5070 3283 270.1189 1.71 47694 57820 251635 12690 584752 30845 20825 274.2736 15.32 73172 7482 73860 75945 74789 7505 7377 275.1253 1.50 130287 320607 48873 36275 1332334 13956 126400 282.8828 1.35 438 589 394 370686 1403 148 1956 296.1212 8.21 1314 17654 686 380718 1428 388 1926 31			424	667	372	375514	1603	155	176
269.1236 6.38 44850 45283 2195 538933 3111 607 60111 269.1241 3.76 9435 76840 2128 1098940 5416 726 3938 269.1241 5.35 5368 36910 1708 47269 4103 507 3233 270.1189 1.71 476984 57820 251635 12659 584752 30845 22082 274.2736 15.32 73172 77482 73680 77945 133234 13956 126402 282.8828 1.35 438 589 394 370686 1403 148 196 284.1340 1.90 336839 39988 157867 13967 320152 8342 17795 296.1212 8.21 1314 17654 686 380718 1428 388 19267 311.1458 1.48 179368 33536 26045 4827 769266 5518 26044			2662	1649	1881	1145	3661	97539	2443
269.1241 3.76 9435 76840 2128 108840 5416 726 3958 269.1241 5.35 5368 36910 1708 47769 4103 507 3283 270.1189 1.71 476984 57820 251635 12659 584752 30845 20825 274.2736 15.32 73172 77482 73800 75945 74789 75705 7337 275.1253 1.50 160287 320607 848873 36275 133234 13956 126402 282.8828 1.35 438 589 35767 330052 84827 13623 4108 1966 284.1340 1.90 336839 35968 157867 13967 330152 8422 1314 17654 666 380718 1428 388 1996 311.1458 1.48 179388 157869 155744 153345 114290 446401 25370 350.1566 1.41			4850	45283	2195	535933	3111	607	60119
270.1189 1.71 476984 57820 251635 12659 584752 30845 20825 274.2736 15.32 73172 73482 7360 75945 74789 75705 7397 275.1253 1.50 180287 320607 848873 36275 133234 13956 12640 282.8828 1.35 438 589 394 370686 1403 148 1956 284.1340 1.90 336839 33968 157867 139678 320152 8342 17395 296.121 8.21 1314 17654 686 380718 428 888 1926 311.1458 1.48 17985 155769 15574 15345 114290 4401 25370 350.1566 1.41 784568 33536 260495 4827 765266 5518 26084 407.1786 1.48 434421 15572 150927 1370 319420 1372 18785 <th>269.1241</th> <th></th> <th>9435</th> <th>76840</th> <th>2128</th> <th>1098840</th> <th>5416</th> <th>726</th> <th>39584</th>	269.1241		9435	76840	2128	1098840	5416	726	39584
270.1189 1.71 476984 57820 251635 12659 584752 30845 20825 274.2736 15.32 73172 73482 7360 75945 74789 75705 7397 275.1253 1.50 180287 320607 848873 36275 133234 13956 12640 282.8828 1.35 438 589 394 370686 1403 148 1956 284.1340 1.90 336839 33968 157867 139678 320152 8342 17395 296.121 8.21 1314 17654 686 380718 428 888 1926 311.1458 1.48 17985 155769 15574 15345 114290 4401 25370 350.1566 1.41 784568 33536 260495 4827 765266 5518 26084 407.1786 1.48 434421 15572 150927 1370 319420 1372 18785 <th></th> <th></th> <th></th> <th></th> <th></th> <th>497269</th> <th></th> <th></th> <th>32830</th>						497269			32830
274.2736 15.32 73172 73482 73860 75945 74789 75755 7397 275.1253 1.50 190287 320607 848873 36275 132334 13956 126402 282.8828 1.35 438 589 394 370686 1403 148 1956 284.1340 1.90 336839 39968 157867 13967 320152 8342 17355 296.1212 8.21 1314 17654 686 380718 1428 388 1926. 311.1458 1.41 784568 33356 260495 4827 769266 5518 26648 407.1786 1.48 434421 15572 150927 1370 31420 1372 18785 410.8026 1.33 172 191 137 4143 325 72328 1990 421.1932 1.56 382467 27989 155837 2216 449483 2512 14921 <tr< th=""><th></th><th></th><th>and the second se</th><th></th><th></th><th></th><th></th><th>8</th><th>208252</th></tr<>			and the second se					8	208252
275.1253 1.50 120287 320607 848873 36275 1132334 13956 126402 282.8828 1.35 438 589 394 370686 403 148 1956 284.1340 1.90 336839 39968 157867 13967 320152 8842 17355 296.1212 8.21 1314 17654 686 380718 14280 388 1926 311.1458 1.48 179388 157867 13967 320152 8842 17355 350.1566 1.41 79388 157869 195744 15387 14280 46401 25370 350.1566 1.41 794568 33356 260455 4827 769266 5518 26644 407.1786 1.48 434421 15572 150927 1370 319420 1372 18785 410.8026 1.33 172 191 137 4143 325 72328 1990 421.1932 1.56 382467 27989 155837 2216 449483 2512 </th <th></th> <th></th> <th>73172</th> <th>73482</th> <th>73860</th> <th>75945</th> <th>74789</th> <th>75705</th> <th>73976</th>			73172	73482	73860	75945	74789	75705	73976
282.8828 1.35 438 589 394 370686 1403 148 196 284.1340 1.90 336839 39968 157867 13967 320152 8842 17859 296.1212 8.21 1314 17654 686 380718 1428 388 1996 311.1458 1.48 179388 157869 195744 15383 1428 388 1926 350.1566 1.41 784568 33356 26054 427 769266 5518 26084 407.1786 1.48 434421 15572 150927 1370 319420 1372 18785 410.8026 1.33 172 191 137 4143 325 72328 1990 421.1932 1.56 382467 27989 153837 2216 449483 2512 14921 478.2160 1.64 268450 14308 100801 671 291750 809 10514	275.1253		1680287	320607	848873	36275	1332334	13956	1264051
284.1340 1.90 336839 33968 157867 13967 320152 8342 17395 296.1212 8.21 1314 17654 686 380718 1428 388 1926 311.1458 1.48 179388 157869 195744 153345 114290 46401 25370 350.1566 1.41 784568 33536 260495 4827 769266 5518 26084 407.1786 1.48 434421 15572 150927 1370 319420 1372 18785 410.8026 1.33 172 191 137 4143 325 72328 190 421.1932 1.56 382467 27989 153837 2216 449483 2512 14324 478.2160 1.64 268450 14308 1000801 671 291750 809 16514 505.8861 1.14 216 83 613655 6882 636 121 218			438	589	394	370686	1403	148	196
296.1212 8.21 1314 17654 686 380718 1428 388 1926 311.1458 1.48 179388 157869 195744 153345 114290 46401 25370 350.1566 1.41 784568 33356 260495 4827 769266 5518 26084 407.1786 1.48 434421 15572 150927 1370 319420 1372 18785 410.8026 1.33 172 191 137 4143 325 72328 190 421.1932 1.56 382467 27989 153837 2216 449483 2512 14921 478.2160 1.64 268450 14308 100801 613655 6882 636 121 218 505.8861 1.14 216 83 613655 6882 636 121 218 514.8905 1.15 771 176 67129 32395 2305 356 714			1000		en .				173959
311.1458 1.48 179388 157869 195744 153345 114290 46401 25370 350.1566 1.41 784568 33536 260495 4827 769266 5518 26084 407.1786 1.48 434421 15572 150927 1370 319420 1372 18785 410.8026 1.33 172 191 137 4143 325 72328 190 421.1932 1.56 382467 27989 153837 2216 449483 2512 14921 478.2160 1.64 268450 14308 100801 611655 6682 636 1212 218 505.8861 1.14 216 83 613655 6682 636 1212 218 514.8905 1.15 771 176 2671729 32395 2305 356 714 515.8902 1.14 185 78 310071 4921 393 97 157	296.1212		1314	17654	686	380718	1428	388	19261
350.1566 1.41 784568 33336 260495 4827 769266 5518 26084 407.1786 1.48 434421 15572 150927 1370 319420 1372 18785 410.8026 1.33 172 191 137 4143 325 72328 190 421.1932 1.56 382467 27989 153837 2216 449483 2512 14921 478.2160 1.64 268450 1408 100801 671 291750 809 1051 505.8861 1.14 216 83 618655 6882 636 121 218 514.8905 1.15 771 176 2671729 32395 2205 356 714 515.8902 1.14 185 78 310071 4921 393 97 157 523.8958 1.14 189 53 623647 6700 498 89 206			873	157869	195744	and the second se	114290	46401	253705
407.1786 1.48 434421 15572 150927 1370 319420 1372 15785 410.8026 1.33 172 191 137 4143 325 72328 190 421.1932 1.56 382467 27989 153837 2216 449483 2512 14921 478.2160 1.64 268450 14308 100815 6882 6365 121 21154 514.8905 1.14 216 83 613655 6882 636 121 218 515.8902 1.14 185 78 31071 4921 393 97 157 523.8958 1.14 189 53 62847 6700 498 89 206									260848
410.8026 1.33 172 191 137 4143 325 72328 190 421.1932 1.56 382467 27989 153837 2216 449483 2512 14921 478.2160 1.64 268450 14308 100801 671 291750 809 10514 505.8861 1.14 216 83 613655 6682 636 121 218 514.8905 1.15 771 176 701729 32395 2305 356 714 515.8902 1.14 185 78 310071 4921 393 97 157 523.8958 1.14 189 53 623647 6700 498 89 206						-		-	187852
421.1932 1.56 382467 27989 153837 2216 449483 2512 14921 478.2160 1.64 268450 14308 100801 671 291750 809 10514 505.8861 1.14 216 83 613655 6682 636 121 218 514.8905 1.15 771 176 201729 32395 2305 356 714 515.8902 1.14 185 78 310071 4921 333 97 157 523.8958 1.14 189 53 623647 6700 498 89 206						4143		8001	190
478.2160 1.64 268450 14308 100801 671 291750 809 10514 505.8861 1.14 216 83 613655 6682 636 121 218 514.8905 1.15 771 176 267129 32395 2305 356 714 515.8902 1.14 185 78 310071 4921 393 97 157 523.8958 1.14 189 53 623647 6700 498 89 206					-	-			149212
505.8861 1.14 216 83 613655 6682 636 121 218 514.8905 1.15 771 176 207129 32395 2305 356 714 515.8902 1.14 185 78 310071 4921 393 97 157 523.8958 1.14 189 53 623647 6700 498 89 206			100	-					105140
514.8905 1.15 771 176 167129 32395 2305 356 714 515.8902 1.14 185 78 310071 4921 393 97 157 523.8958 1.14 189 53 623647 6700 498 89 206				-					-
515.8902 1.14 185 78 310071 4921 393 97 157 523.8958 1.14 189 53 623647 6700 498 89 206									
523.8958 1.14 189 53 6 23647 6700 498 89 206						-			
						-			
532.9017 1.14 280 89 578900 8100 669 104 231									

Figure S15. Table of selected features ordered by m/z from the experiments where the mineral present was varied. Features were selected from a full list based on absolute MS intensity (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	Alumina	Montmoril.	Mica	Goethite	Quartz	Natrolite	Silica
515.8902	1.14	185	78	310071	4921	393	97	157
505.8861	1.14	216	83	613655	6882	636	121	218
528.8886	1.14	162	91	334769	4334	382	61	152
523.8958	1.14	189	53	623647	6700	498	89	206
532.9017	1.14	280	89	578900	8100	669	104	231
514.8905	1.15	771	176	2671729	32395	2305	356	714
182.9022	1.33	1405	2779	40968	1180	77493	938129	13335
410.8026	1.33	172	191	137	4143	325	72328	190
154.9073	1.33	337	441	6695	210	12237	140056	2254
181.9025	1.33	562	887	6174	2370	11727	137192	2148
180.9015	1.33	255	430	3994	2566	7377	91702	1405
179.9021	1.33	222	349	3436	1159	6429	76960	1260
215.6010	1.33	304632	5532	92751	1542	336540	2383	138847
244.1118	1.34	381586	6170	104192	3535	410070	1199	136826
282.8828	1.35	438	589	394	370686	1403	148	196
246.8617	1.36	399	872	419	432553	2090	352	244
264.8731	1.36	424	667	372	375514	1603	155	176
152.5797	1.40	357541	16051	134391	3551	341267	6916	128251
147.0716	1.40	740358	51620	398704	10448	749182	9076	350371
350.1566	1.41	784568	33536	260495	4827	769266	5518	260848
168.0768	1.44	19697	345645	31007	453617	19130	24846	117762
95.0607	1.47	145157	289066	138974	342833	146885	38298	234199
175.5824	1.48	546595	27370	201717	5460	404957	7829	213271
311.1458	1.48	179388	157869	195744	153345	114290	46401	253705
111.0742	1.48	225188	77487	150542	79025	241627	39939	165199
407.1786	1.48	434421	15572	150927	1370	319420	1372	187852
182.0917	1.49	13042	169030	21222	201906	24374	17222	27646
275.1253	1.50	1680287	320607	848873	36275	1332334	13956	1264051
267.1082	1.53	2662	1649	1881	1145	3661	97539	2443
421.1932	1.56	382467	27989	153837	2216	449483	2512	149212
156.0765	1.59	2024089	724691	1831942	1023374	2309911	419797	1880253
175.0974	1.63	81035	247721	242281	48134	42781	20891	244586
167.0925	1.64	173668	138856	220184	59014	198455	7332	91836
478.2160	1.64	268450	14308	100801	671	291750	809	105140
157.0793	1.65	155643	64398	114414	66553	150392	29600	122255
248.1140	1.66	90834	509572	519967	287227	72684	14117	1276975
227.1133	1.67	379633	87850	175657	53876	358354	25397	183543
270.1189	1.71	476984	57820	251635	12659	584752	30845	208252
211.1010	1.72	219607	14722	77880	18768	339656	4001	102547
NUCLEON DESCRIPTION	1.72	338982	14722	101541	2438	284942	1909	139587
239.6115		1011				1000 C	and a second sec	
213.0977	1.78	311076	85252	166383	58108	121052	113360	351759
196.0901	1.80	287575	163615	231349	31968	163565	3560	457434
195.0876	1.80	1560203	2007331	2897008	390669	2019202	34010	5572833
110.0715	1.81	1035866	687981	446650	142986	1783821	131242	1759590
284.1340	1.90	336839	39968	157867	13967	320152	8342	173959
184.0717	1.98	32902	214129	21188	352553	33487	30513	119416
241.0928	2.05	21817	132535	18589	174107	27104	4429	90458
181.1078	2.08	300721	152646	225367	36155	175437	2490	449602
209.1030	2.22	893142	401418	2971439	380200	1969699	28795	8189056
210.1054	2.23	348576	181170	275068	40840	195902	3370	552761
209.0669	2.29	1291658	694515	1016808	119483	689164	11990	878018
227.0769	2.30	29228	49306	16908	200786	15244	1583	248325
180.0765	2.95	10720	71250	3826	660407	7113	1453	38173
198.0868	3.30	6111	115601	4644	141804	4203	612	56031
269.1241	3.76	9435	76840	2128	1098840	5416	726	39584
156.0767	3.77	21543	62783	7596	574864	22279	1512	69405
211.0686	4.66	1599	26368	941	342118	1604	409	9850
180.0763	5.35	3083	26431	941	343724	2102	504	19676
269.1241	5.35	5368	36910	1708	497269	4103	507	32830
180.0764	6.38	4334	44252	1767	424188	2994	463	47418
269.1236	6.38	4850	45283	2195	535933	3111	607	60119
156.0764	7.40	185999	99617	67575	121150	248873	12003	664988
110.0713	7.41	364164	351148	84117	568845	435159	11725	958289
296.1212	8.21	1314	17654	686	380718	1428	388	19261
246.2419	13.84	34852	34129	33625	33828	33949	34868	32923
274.2736	15.34	73172	73482	73860	75945	74789	75705	73976

Figure S16. Table of selected features ordered by RT from the experiments where the mineral present was varied. Features were selected from a full list based on absolute MS intensity (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	Alumina	Montmoril.	Mica	Goethite	Quartz	Natrolite	Silica
190.0821	106.33	81845	14625	64412	2696	63245	663	28069
204.0917	99.46	219778	11223	91490	21652	261853	3306	97548
204.0962	127.55	60094	19200	46597	17070	52550	697	47758
218.1088	109.19	55221	65487	23351	55581	63822	1646	35161
218.1124	165.03	43244	23870	29818	10113	35623	414	40421
261.1164	107.09	78025	16103	38199	3676	94540	851	40628
261.1189	130.35	48288	14500	31787	2333	52898	326	31099
270.1166	446.33	12546	10604	4614	22820	12965	739	37557
270.1189	102.58	476984	57820	251635	12659	584752	90845	208252
270.1193	203.53	4498	3642	1570	11178	4786	190	5796
270.1253	437.36	7112	7826	2475	22493	8374	302	13584
270.1260	382.91	1335	4881	498	73487	1266	174	8236
270.1262	225.49	2809	10693	969	144392	2762	182	8725
270.1265	320.95	1393	5365	613	69264	1364	200	5434
275.1253	90.10	680287	920607	948873	36275	1382384	18956	264051
275.1331	160.07	76346	25521	43215	5518	80817	465	56189
284.1340	114.07	336839	39968	157867	13967	320152	8342	173959
298.1506	121.59	63768	11648	64090	10665	54499	909	84583
318.1343	103.55	32158	5603	10901	4434	33418	738	12331
327.1409	105.98	178029	25541	96241	1735	189785	3062	97626
327.1484	302.93	1444	9041	546	40870	1214	126	5067
327.1486	368.38	1657	7770	602	46555	1385	203	3226
332.1463	81.18	80929	4818	18979	2747	102345	876	35578
341.1558	435.57	10500	4917	2327	13382	11155	101	7557
341.1559	129.56	221120	24203	90918	4794	220460	3278	98063
341.1613	455.36	3939	11639	836	37175	2328	150	7381
341.1642	380.41	1520	5570	551	33667	1799	116	5046
341.1642	455.34	4847	11560	1037	39941	2998	189	10215
350.1566	84.62	784568	33536	260495	4827	769266	5518	260848
355.1716	125.59	105409	16154	51034	3063	97258	496	51696
364.1713	88.05	265593	25441	128196	6017	219282	3101	111938
384.1613	111.56	86028	20459	45497	72	119769	356	43594
398.1759	126.59	80955	22512	52355	1015	77728	363	56508
398.1835	454.95	3730	18770	1417	31025	4187	199	11083
407.1786	89.09	434421	15572	150927	1370	319420	1372	187852
412.1975	463.38	873	3593	347	14039	920	108	2437
421.1932	93.83	382467	27989	153837	2216	449483	2512	149212
441.1834	102.61	137738	10334	74373	578	144104	458	67026
455.1984	124.55	117057	9560	41989	409	109966	176	41297
464.1984	91.94	170895	8581	81254	677	187251	876	82476
478.2160	98.11	268450	14308	100801	671	291750	809	105140
487.2164	80.10	93232	840	14317	324	93229	218	38993
492.2312	105.05	66624	6390	36886	734	109139	303	36040
535.2355	103.62	164530	6789	46567	538	175398	338	59685

Figure S17. Table of selected features ordered by m/z from the experiments where the mineral present was varied. Features were selected from a full list based on absolute MS intensity, filtered to include masses consistent with peptides, (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	Alumina	Montmoril.	Mica	Goethite	Quartz	Natrolite	Silica
487.2164	80.10	93232	840	14317	324	93229	218	38993
332.1463	81.18	80929	4818	18979	2747	102345	876	35578
350.1566	84.62	784568	33536	260495	4827	769266	5518	260848
364.1713	88.05	265593	25441	128196	6017	219282	3101	111938
407.1786	89.09	434421	15572	150927	1370	319420	1372	187852
275.1253	90.10	680287	120607	848873	36275	1332334	18956	1264051
464.1984	91.94	170895	8581	81254	677	187251	876	82476
421.1932	93.83	382467	27989	153837	2216	449483	2512	149212
478.2160	98.11	268450	14308	100801	671	291750	809	105140
204.0917	99.46	219778	11223	91490	21652	261853	3306	97548
270.1189	102.58	476984	57820	251635	12659	584752	10845	208252
441.1834	102.61	137738	10334	74373	578	144104	458	67026
318.1343	103.55	32158	5603	10901	4434	33418	738	12331
535.2355	103.62	164530	6789	46567	538	175398	338	59685
492.2312	105.05	66624	6390	36886	734	109139	303	36040
327.1409	105.98	178029	25541	96241	1735	189785	3062	97626
190.0821	106.33	81845	14625	64412	2696	63245	663	28069
261.1164	107.09	78025	16103	38199	3676	94540	851	40628
218.1088	109.19	55221	65487	23351	55581	63822	1646	35161
384.1613	111.56	86028	20459	45497	72	119769	356	43594
284.1340	114.07	336839	39968	157867	13967	320152	8342	173959
298.1506	121.59	63768	11648	64090	10665	54499	909	84583
455.1984	124.55	117057	9560	41989	409	109966	176	41297
355.1716	125.59	105409	16154	51034	3063	97258	496	51696
398.1759	126.59	80955	22512	52355	1015	77728	363	56508
204.0962	127.55	60094	19200	46597	17070	52550	697	47758
341.1559	129.56	221120	24203	90918	4794	220460	3278	98063
261.1189	130.35	48288	14500	31787	2333	52898	326	31099
275.1331	160.07	76346	25521	43215	5518	80817	465	56189
218.1124	165.03	43244	23870	29818	10113	35623	414	40421
270.1193	203.53	4498	3642	1570	11178	4786	190	5796
270.1262	225.49	2809	10693	969	144392	2762	182	8725
327.1484	302.93	1444	9041	546	40870	1214	126	5067
270.1265	320.95	1393	5365	613	69264	1364	200	5434
327.1486	368.38	1657	7770	602	46555	1385	203	3226
341.1642	380.41	1520	5570	551	33667	1799	116	5046
270.1260	382.91	1335	4881	498	73487	1266	174	8236
341.1558	435.57	10500	4917	2327	13382	11155	101	7557
270.1253	437.36	7112	7826	2475	22493	8374	302	13584
270.1166	446.33	12546	10604	4614	22820	12965	739	37557
398.1835	454.95	3730	18770	1417	31025	4187	199	11083
341.1642	455.34	4847	11560	1037	39941	2998	189	10215
341.1613	455.36	3939	11639	836	37175	2328	150	7381
412.1975	463.38	873	3593	347	14039	920	108	2437

Figure S18. Table of selected features ordered by RT from the experiments where the mineral present was varied. Features were selected from a full list based on absolute MS intensity, filtered to include masses consistent with peptides, (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	G>A>H	G>H>A	A > G > H	A > H > G	H > G > A	H > A > G	G+A+H
110.0714	2.89	471128	543529	536493	265449	451612	272511	645694
110.0714	7.35	667261	838782	654055	471790	435025	398691	1038814
110.0716	1.40	4310865	12937166	4929553	14183757	11389903	15484259	9653760
115.0503	2.09	838965	682842	432764	159365	390690	216411	405412
124.0691	1.38	427297	622792	351939	525940	629125	682310	480923
147.0721	1.42	1053929	1307296	1135616	861143	965312	694040	930390
147.0762	2.32	474473	376912	704564	268544	415692	348935	526425
152.5798	1.42	255707	572162	242183	271360	626590	233528	576079
156.0768	1.49	1732736	3069003	#133024	2393722	3222443	2499476	4856198
167.0926	1.79	425341	802161	208834	223789	623431	309748	494064
175.5825	1.62	152386	946728	478523	518423	1113972	287992	507549
181.1081	2.20	522636	297973	119726	318220	100901	501177	256897
182.5903	1.72	532747	355364	261112	467896	276846	630746	571794
195.0876	1.80	2364110	3146026	664353	966257	1695550	1637330	2154442
209.1032	2.21	8976109	2965291	844627	2552330	702686	4923785	2246411
211.1012	1.78	542947	618478	454928	462284	486232	450113	757126
213.0982	1.85	559593	524542	809297	307778	577368	330075	518951
215.6014	1.34	1071839	1375281	987155	1667987	1125751	2014942	1068491
239.6117	2.10	529095	567038	875981	111648	113484	101609	531090
244.1122	1.35	575542	1567016	631913	611350	1418008	628841	1242060
247.1299	1.38	241005	414426	216241	453906	384545	548599	329979
251.1193	1.38	301352	198747	269873	1089853	153931	1090442	351229
268.1224	1.97	278874	680959	291488	351853	813757	211222	629518
274.2742	15.33	994574	725904	721863	572348	638025	799601	739803
275.1255	1.51	1298691	1327991	1395821	1906366	1297633	2696740	853237
284.1338	2.03	229407	517629	565688	299994	686321	437834	523436
284.1359	2.13	572644	473667	907752	517481	482644	548557	393422
319.6496	1.28	76304	92624	52832	429097	74009	506950	148239
327.1410	1.81	710154	557269	406481	250910	262924	144435	521404
341.1570	2.17	1275800	871469	1331522	385257	654204	403212	807927
350.1572	1.61	1127638	2487113	908794	911294	2544588	1009937	2071712
398.1777	1.91	3053892	757155	2480547	785277	1069829	475137	639337
399.1802	2.09	582613	88955	441184	144940	52256	62777	199140
407.1786	1.50	452036	1034357	549936	396684	1011150	314605	967019
412.1932	2.32	627135	153603	400534	140334 846818	106363 492980	163610 363513	300995 745658
421.1929	1.61	581778	586913	Charles Contraction	531549	492980	787063	
430.1946	1.31	372267	551536	296702 132560	903633	85607	688989	317248
435.2102	1.76	1350796	Construction of the	1096215	247684	86830	208250	705904
455.1968 455.2000	2.05	933996	330202 339872	734942	139844	275646	136322	570742
455.2000	1.53	187840	623974	343493	148254	596959	96734	361115
478.2153	1.53	588171	506087	647919	433932	401571	362650	706492
478.2153	1.34	322126	1227204	305573	364005	1029196	415941	871706
501.2325	1.34	226611	169836	169664	898684	118565	924449	290690
535.2378	1.83	513340	553150	548248	224086	469476	184038	718262
555.2578	1.83	513340	000000	548248	224086	403475	184038	/18262

Figure S19. Table of selected features ordered by m/z from the experiments where the order of addition was varied. Features were selected from a full list based on absolute MS intensity (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	G > A > H	G > H > A	A > G > H	A > H > G	H > G > A	H > A > G	G+A+H
319.6496	1.28	76304	92624	52832	429097	74009	506950	148239
430.1946	1.31	372267	551536	296702	531549	441449	787063	317248
487.2154	1.34	322126	1227204	305573	364005	1029196	415941	871706
215.6014	1.34	1071839	1375281	987155	1667987	1125751	2014942	1068491
244.1122	1.35	575542	1567016	631913	611350	1418008	628841	1242060
501.2325	1.37	226611	169836	169664	898684	118565	924449	290690
251.1193	1.38	301352	198747	269873	1089853	153931	1090442	351229
124.0691	1.38	427297	622792	351939	525940	629125	682310	480923
247.1299	1.38	241005	414426	216241	453906	384545	548599	329979
110.0716	1.40	4310865	12937166	4929553	(#183757	1389903	15484259	1653760
152.5798	1.42	255707	572162	242183	271360	626590	233528	576079
147.0721	1.42	1053929	1307296	1135616	861143	965312	694040	930390
156.0768	1.49	1732736	3069003	9133024	2393722	3222443	2499476	4856198
407.1786	1.50	452036	1034357	549936	396684	1011150	314605	967019
275.1255	1.51	1298691	1327991	1395821	1906366	1297633	2696740	853237
464.2003	1.53	187840	623974	343493	148254	596959	96734	361115
350.1572	1.61	1127638	2487113	908794	911294	2544588	1009937	2071712
421.1929	1.61	581778	586913	762372	846818	492980	363513	745658
175.5825	1.62	152386	946728	478523	518423	1113972	287992	507549
182.5903	1.72	532747	355364	261112	467896	276846	630746	571794
435.2102	1.76	180496	165802	132560	903633	85607	688989	185113
211.1012	1.78	542947	618478	454928	462284	486232	450113	757126
167.0926	1.79	425341	802161	208834	223789	623431	309748	494064
195.0876	1.80	2864110	3146026	664353	966257	1695550	1637330	2154442
327,1410	1.81	710154	557269	406481	250910	262924	144435	521404
535.2378	1.83	513340	553150	548248	224086	469476	184038	718262
213.0982	1.85	559593	524542	809297	307778	577368	330075	518951
478,2153	1.86	588171	506087	647919	433932	401571	362650	706492
398.1777	1.91	3053892	757155	2480547	785277	1069829	475137	639337
268.1224	1.97	278874	680959	291488	351853	813757	211222	629518
284.1338	2.03	229407	517629	565688	299994	686321	437834	523436
455.1968	2.05	1350796	330202	1096215	247684	86830	208250	705904
115.0503	2.09	838965	682842	432764	159365	390690	216411	405412
399.1802	2.09	582613	88955	441184	144940	52256	62777	199140
239.6117	2.10	529095	567038	875981	111648	113484	101609	531090
284.1359	2.13	572644	473667	907752	517481	482644	548557	393422
341.1570	2.17	1275800	871469	1331522	385257	654204	403212	807927
181,1081	2.20	522636	297973	119726	318220	100901	501177	256897
209.1032	2.21	5976109	2965291	844627	2552330	702686	4923785	2246411
455.2000	2.23	933996	339872	734942	139844	275646	136322	570742
147.0762	2.32	474473	376912	704564	268544	415692	348935	526425
412.1932	2.32	627135	153603	400534	140334	106363	163610	300995
110.0714	2.89	471128	543529	536493	265449	451612	272511	645694
110.0714	7.35	667261	838782	654055	471790	435025	398691	1038814
274.2742	15.33	994574	725904	721863	572348	638025	799601	739803

Figure S20. Table of selected features ordered by RT from the experiments where the order of addition was varied. Features were selected from a full list based on absolute MS intensity (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	G>A>H	G>H>A	A > G > H	A>H>G	H>G>A	H > A > G	G + A + H
261.1095	14.99	6654	8580	5712	5250	8018	5567	6173
303.1659	4.35	4360	7439	4387	4066	3351	4617	2466
318.1495	1.26	4647	40085	3887	9022	32531	9227	23708
360.1874	6.20	6843	1481	11505	1879	834	1941	2495
369.1870	7.37	13644	1012	14550	7117	173	6905	2841
389.1801	1.24	5127	28436	5235	7555	25747	10203	20515
421.1879	1.20	782	5375	476	3523	7016	10662	4384
441.1834	3.77	6949	2457	11036	815	1020	735	2331
503.2205	6.67	17922	4023	9234	445	1165	370	1171
517.2361	5.45	21153	6037	21562	721	2450	596	2620
526.2279	1.29	6176	31774	5526	4980	25873	5090	19777
535.2359	7.24	2254	3177	4251	1027	2199	758	6080
540.2512	7.39	26052	3092	64913	2602	1034	2425	8300
540.2524	3.00	9392	1181	11635	681	364	604	2096
540.2530	3.82	12129	2249	22842	1237	1005	931	4551
549.2433	1.53	15118	10092	3363	11002	4239	35536	3506
560.2398	7.31	11881	3317	6207	530	897	454	1551
560.2424	4.45	43602	11237	28659	578	2981	500	2717
589.2328	2.15	15498	11479	17212	917	4321	685	2504
603.2478	3.29	49740	4405	35296	618	1932	482	3016
603.2479	2.54	10324	4333	14188	510	1603	489	1605
617.2628	7.26	22665	3677	18210	487	1172	463	1999
617.2642	4.24	8505	874	12216	236	425	165	1234
625.3039	7.55	7465	874	14754	1557	424	1305	3223
631.2796	7.54	56211	4217	78758	595	1678	501	5945
640.2796	2.68	19249	4022	13964	594	1183	458	2770
643.3019	1.70	2265	1637	2384	18354	744	13586	4647
645.2954	7.72	19605	1307	36882	449	513	355	3043
668.3101	7.46	21435	3018	60204	1578	1665	1228	9722
674.2857	7.35	19007	1828	12010	493	531	361	1301
682.3268	7.58	5430	577	12287	587	381	489	2030
683.2855	2.67	47599	18490	20066	512	2814	397	3264
688.3007	7.52	26192	1836	34844	450	530	357	2146
695.3210	1.82	6657	2889	7200	6008	1670	7735	4774
702.3161	7.70	11769	550	19995	295	448	241	1309
704.3151	1.23	3521	9112	941	6802	9743	18000	6582
709.3290	1.38	4274	4901	3611	37422	3219	38958	10926
715.3107	1.99	5135	9036	5957	817	2154	730	4517
725.3310	7.47	12126	1947	36525	729	850	615	4449
749.3149	2.41	30671	22498	9519	522	4333	388	1566
761.3324	1.22	958	17738	650	3889	17126	10013	8537
763.3274	2.66	11333	5995	5603	336	2317	231	3309
768.3379	7.38	15373	2268	27225	474	1007	359	3749
772.3250	1.63	7021	30100	3258	1611	22554	1150	11415
775.3511	1.24	678	939	140	6997	1100	15091	1772
782.3533	7.52	7457	419	18283	287	292	281	1571
795.3382	1.38	7345	54638	5588	5234	50632	5188	30129
809.3562	1.45	2896	10600	4756	5111	9392	4598	9731
818.3554	1.24	659	19509	567	1813	13690	4682	8101
829.3456	1.68	3187	18174	2060	702	13259	584	7975
841.3808	1.53	6471	4977	14277	13589	4687	9460	11581
852.3630	1.43	6374	18296	2588	4935	17179	5556	10408
875.3790	1.25	620	15059	455	959	11457	1594	9515

Figure S21. Table of selected features ordered by m/z from the experiments where the order of addition was varied. Features were selected from a full list based on absolute MS intensity, filtered to include masses consistent with peptides, (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	G > A > H	G > H > A	A > G > H	A > H > G	H > G > A	H > A > G	G+A+H
421.1879	1.20	782	5375	476	3523	7016	10662	4384
761.3324	1.22	958	17738	650	3889	17126	10013	8537
704.3151	1.23	3521	9112	941	6802	9743	18000	6582
389.1801	1.24	5127	28436	5235	7555	25747	10203	20515
818.3554	1.24	659	19509	567	1813	13690	4682	8101
775.3511	1.24	678	939	140	6997	1100	15091	1772
875.3790	1.25	620	15059	455	959	11457	1594	9515
318.1495	1.26	4647	40085	3887	9022	32531	9227	23708
526.2279	1.29	6176	31774	5526	4980	25873	5090	19777
795.3382	1.38	7345	54638	5588	5234	50632	5188	30129
709.3290	1.38	4274	4901	3611	37422	3219	33958	10926
852.3630	1.43	6374	18296	2588	4935	17179	5556	10408
809.3562	1.45	2896	10600	4756	5111	9392	4598	9731
841.3808	1.53	6471	4977	14277	13589	4687	9460	11581
549.2433	1.53	15118	10092	3363	11002	4239	35536	3506
772.3250	1.63	7021	30100	3258	1611	22554	1150	11415
829.3456	1.68	3187	18174	2060	702	13259	584	7975
643.3019	1.70	2265	1637	2384	18854	744	13586	4647
695.3210	1.82	6657	2889	7200	6008	1670	7735	4774
715.3107	1.99	5135	9036	5957	817	2154	730	4517
589.2328	2.15	15498	11479	17212	917	4321	685	2504
749.3149	2.41	30671	22498	9519	522	4333	388	1566
603.2479	2.54	10324	4333	14188	510	1603	489	1605
763.3274	2.66	11333	5995	5603	336	2317	231	3309
683.2855	2.67	47599	18490	20066	512	2814	397	3264
640.2796	2.68	19249	4022	13964	594	1183	458	2770
540.2524	3.00	9392	1181	11635	681	364	604	2096
603.2478	3.29	49740	4405	35296	618	1932	482	3016
441.1834	3.77	6949	2457	11036	815	1020	735	2331
540,2530	3.82	12129	2249	22842	1237	1005	931	4551
617.2642	4.24	8505	874	12216	236	425	165	1234
303.1659	4.35	4360	7439	4387	4066	3351	4617	2466
560.2424	4.45	43602	11237	28659	578	2981	500	2717
517.2361	5.45	21153	6037	21562	721	2450	596	2620
360.1874	6.20	6843	1481	11505	1879	834	1941	2495
503.2205	6.67	17922	4023	9234	445	1165	370	1171
535.2359	7.24	2254	3177	4251	1027	2199	758	6080
617.2628	7.26	22665	3677	18210	487	1172	463	1999
560.2398	7.31	11881	3317	6207	530	897	454	1551
674.2857	7.35	19007	1828	12010	493	531	361	1301
369.1870	7.37	13644	1012	14550	7117	173	6905	2841
768.3379	7.38	15373	2268	27225	474	1007	359	3749
540.2512	7.39	26052	3092	64913	2602	1034	2425	8300
668.3101	7.46	21435	3018	60204	1578	1665	1228	9722
725.3310	7.47	12126	1947	36525	729	850	615	4449
782.3533	7.52	7457	419	18283	287	292	281	1571
688.3007	7.52	20192	1836	34844	450	530	357	2146
631.2796	7.54	56211	4217	78758	595	1678	501	5945
625.3039	7.55	7465	874	14754	1557	424	1305	3223
682.3268	7.58	5430	577	12287	587	381	489	2030
702.3161	7.70	11769	550	19995	295	448	241	1309
645.2954	7.72	19605	1307	36882	449	513	355	3043
261.1095	14.99	6654	8580	5712	5250	8018	5567	6173

Figure S22. Table of selected features ordered by RT from the experiments where the order of addition was varied. Features were selected from a full list based on absolute MS intensity, filtered to include masses consistent with peptides, (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

Picked m/z	RT (min)	Int (avg)	Matched m/z	Difference	Formula
204.0953	1.67	466368	204.0977	-0.0024	G2A
218.1134	2.62	<mark>29046</mark>	218.1133	0.0000	GA2
261.1160	1.78	86010	261.1192	-0.0032	G3A
270.1215	1.67	442474	270.1195	0.0020	G2H
275.1326	2.66	62228	275.1349	-0.0022	G2A2
284.1327	1.91	341876	284.1351	-0.0025	GAH
298.1533	2.25	66118	298.1508	0.0025	A2H
327.1430	1.78	188869	327.1410	0.0020	G3H
341.1565	2.17	160775	341.1567	-0.0001	G2AH
350.1550	1.41	1267554	350.1569	-0.0019	GH2
355.1702	1.99	112959	355.1723	-0.0021	GA2H
364.1755	1.50	615381	364.1726	0.0029	AH2
369.1904	2.78	18881	369.1880	0.0025	A3H
384.1650	1.64	234681	384.1625	0.0025	G4H
398.1789	1.89	199310	398.1782	0.0008	G3AH
407.1804	1.46	478800	407.1785	0.0019	G2H2
421.1984	1.58	612085	421.1941	0.0043	GAH2
430.1945	1.37	147853	430.1944	0.0001	H3
435.2125	1.70	110105	435.2098	0.0028	A2H2
487.2180	1.33	278866	487.2159	0.0021	GH3
567.2564	1.24	15493	567.2533	0.0030	H4

Figure S23. Table of consistent peptide compositions for selected features from the experiments where the salts present were varied. Selected features were filtered to include masses consistent with peptides of 3- to 15-mer of G, A and H, and the list of features further refined by selecting only those that appeared in the top 20 (ranked by absolute intensity) for at least one condition in the experiment. Intensity for each feature is in absolute counts (averaged over experimental and analytical replicates). Note: these compositions are only consistent with the mass observed, but no further validation has been carried out. Furthermore, without sequence information little or no conclusion can be drawn from the formulae – see following section.

Picked m/z	RT (min)	Int (avg)	Matched m/z	Difference	Formula
190.0821	1.77	81845	190.0820	0.0001	G3
204.0962	2.13	60094	204.0977	-0.0015	G2A
218.1124	2.75	43244	218.1133	-0.0010	GA2
261.1189	2.17	52898	261.1192	-0.0003	G3A
270.1193	3.39	11178	270.1195	-0.0002	G2H
275.1331	2.67	80817	275.1349	-0.0017	G2A2
284.1340	1.90	336839	284.1351	-0.0012	GAH
298.1506	2.03	84583	298.1508	-0.0002	A2H
327.1409	1.77	189785	327.1410	-0.0001	G3H
341.1559	2.16	221120	341.1567	-0.0007	G2AH
350.1566	1.41	784568	350.1569	-0.0003	GH2
355.1716	2.09	105409	355.1723	-0.0007	GA2H
364.1713	1.47	265593	364.1726	-0.0013	AH2
384.1613	1.86	119769	384.1625	-0.0012	G4H
398.1759	2.11	80955	398.1782	-0.0023	G3AH
407.1786	1.48	434421	407.1785	0.0001	G2H2
412.1975	7.72	14039	412.1938	0.0037	G2A2H
421.1932	1.56	449483	421.1941	-0.0009	GAH2
441.1834	1.71	144104	441.1840	-0.0006	G5H
455.1984	2.08	117057	455.1997	-0.0013	G4AH
464.1984	1.53	187251	464.2000	-0.0016	G3H2
478.2160	1.64	291750	478.2156	0.0004	G2AH2
487.2164	1.33	93232	487.2159	0.0005	GH3
492.2312	1.75	109139	492.2313	0.0000	GA2H2
535.2355	1.73	175398	535.2371	-0.0016	G3AH2

Figure S24. Table of consistent peptide compositions for selected features from the experiments where the minerals present were varied. Selected features were filtered to include masses consistent with peptides of 3- to 15-mer of G, A and H, and the list of features further refined by selecting only those that appeared in the top 20 (ranked by absolute intensity) for at least one condition in the experiment. Intensity for each feature is in absolute counts (averaged over experimental and analytical replicates). Note: these compositions are only consistent with the mass observed, but no further validation has been carried out. Furthermore, without sequence information little or no conclusion can be drawn from the formulae – see following section.

Picked m/z	RT (min)	Int (avg)	Matched m/z	Difference	Formula
303.1659	4.35	7439	303.1662	-0.0002	A4
360.1874	6.20	11505	360.1877	-0.0003	GA4
369.1870	7.37	14550	369.1880	-0.0009	A3H
389.1801	1.24	28436	389.1779	0.0023	G4A2
441.1834	3.77	11036	441.1840	-0.0006	G5H
503.2205	6.67	17922	503.2209	-0.0004	G6A2
517.2361	5.45	21562	517.2365	-0.0005	G5A3
535.2359	7.24	6080	535.2371	-0.0012	G3AH2
540.2524	3.00	11635	540.2525	-0.0001	G3A3H
560.2424	4.45	43602	560.2424	0.0000	G7A2
589.2328	2.15	17212	589.2326	0.0002	G10
603.2479	2.54	14188	603.2483	-0.0004	G9A
617.2642	4.24	12216	617.2639	0.0003	G8A2
625.3039	7.55	14754	625.3053	-0.0014	G2A5H
631.2796	7.54	78758	631.2796	0.0000	G7A3
640.2796	2.68	1 9249	640.2798	-0.0003	G6A2H
643.3019	1.70	18354	643.3059	-0.0040	A3H3
645.2954	7.72	36882	645.2952	0.0002	G6A4
668.3101	7.46	60204	668.3111	-0.0011	G4A4H
674.2857	7.35	19007	674.2854	0.0003	G9A2
682.3268	7.58	12287	682.3268	0.0000	G3A5H
683.2855	2.67	47599	683.2857	-0.0002	G8AH
688.3007	7.52	34844	688.3011	-0.0003	G8A3
702.3161	7.70	19995	702.3167	-0.0006	G7A4
704.3151	1.23	18000	704.3123	0.0028	H5
709.3290	1.38	37422	709.3277	0.0013	A2H4
725.3310	7.47	36525	725.3327	-0.0016	G5A4H
761.3324	1.22	17738	761.3338	-0.0014	GH5
763.3274	2.66	11333	763.3232	0.0042	G7AH2
768.3379	7.38	27225	768.3385	-0.0006	G7A3H
772.3250	1.63	30100	772.3234	0.0015	G6H3
775.3511	1.24	15091	775.3495	0.0017	AH5
782.3533	7.52	18283	782.3542	-0.0009	G6A4H
795.3382	1.38	54638	795.3394	-0.0012	G4H4
809.3562	1.45	10600	809.3550	0.0011	G3AH4
818.3554	1.24	19509	818.3553	0.0001	G2H5
829.3456	1.68	18174	829.3450	0.0006	G7H3
852.3630	1.43	18296	852.3609	0.0021	G5H4
875.3790	1.25	15059	875.3768	0.0021	G3H5

Figure S25. Table of consistent peptide compositions for selected features from the experiments where the order of addition was varied. Selected features were filtered to include masses consistent with peptides of 3- to 15-mer of G, A and H, and the list of features further refined by selecting only those that appeared in the top 20 (ranked by absolute intensity) for at least one condition in the experiment. Intensity for each feature is in absolute counts (averaged over experimental and analytical replicates). Note: these compositions are only consistent with the mass observed, but no further validation has been carried out. Furthermore, without sequence information little or no conclusion can be drawn from the formulae – see following section. Page 37

2.2.3 Sequence permutation distribution difference between populations

As outlined above, our aim in LC-MS analysis was to characterise product distribution without the bias/distraction associated with product expectations. We see clearly, both in populationlevel analyses, and in simple observation of extracted ion chromatograms (EICs) of particular m/z values, that product distribution differs clearly and consistently. Since the (secondary & higher) structure and function of oligomeric species depend not only on their composition (e.g. which AAs are incorporated), but also on the sequence of monomers, it is instructive to ask: 'Is the sequence of oligomer products altered by the conditions being manipulated?".

To answer this question unequivocally is difficult; however, it requires identifying and separating very similar species, including those of identical mass. In many cases such isomeric species are extremely difficult to resolve using chromatography – even more so when the chromatography method is general, rather than optimised to resolve specific sequence variants. Below, we show an example where discrete peaks in chromatograms can be assigned to correspond to particular species and demonstrate that different product ensembles can incorporate different sequence permutation distributions (Figure S26; the basis of these assignments explained in Figure 27.

Further examples of sets of isobaric species (likely isomeric/different sequence permutations in many cases) in which both relative and absolute amounts observed in different ensembles vary markedly can be seen in Figures S8 to S10.




Figure S26. Plots revealing the sequence permutation distribution of G_4A pentamers. (a) EICs of m/z = 318.141 from products of a mixing history experiment; (b) EICs from part (a) normalised to respective maxima; (c) Identification of sequence permutations contributing to each peak; labels colours followed in intensity plots; (d) Distribution of mean intensity for samples of different mixing histories, with error bars representing one standard deviation; (e) Distribution of mean intensity for samples of different mixing histories; peaks intensities were extracted using Bruker Data analysis as integrated intensities following peak picking; intensity values displayed are means of all 9 data sets (3 experimental reps x 3 analytical reps), with error bars representing one standard deviation]




Figure S27. The basis for assignment of the three peaks observed in EICs corresponding to G₄A oligomers. (a) EIC of m/z = 318.1408 +/-0.002 in a G, A, H condensation product ensemble ("Product Mix") compared to base peak chromatograms of standards of the five sequence permutations possible in G₄A pentamers; this comparison confirms the assignment of peak identity shown in (b), and is consistent with data from attempts at de novo assignment using MS^2 . (c) Example MS^2 spectrum, derived from the fragmentation of m/z = 318.1408 with retention time at 2.1 mins, showing fragmentation consistent with assignment as co-elution of GAGGG, GGAGG, and GGGAG pentamers (middle peak in S27b).

Note: It was necessary to use synthetic standards (produced by standard SPPS) to confirm the identity of each peak, as robust unequivocal *de novo* assignment is not possible solely based on MS² data. MS² analysis of each peak did yield fragments consistent with β - and γ - series derived from the sequences finally assigned, however, other peaks were also observed which were consistent with <u>other</u> sequences. For example, MS² spectra of the first peak, which corresponds to AGGGG, included a strong peak with m/z = 151.0502: this is consistent with Simple β - or γ - fragments of the AGGGG sequence. We speculate that this might result from a Maclaffery Rearrangement.

We include this note to illustrate that robust unequivocal *de novo* assignment of abiotic peptide sequence, where many of the possible sequence permutations are present, is not facile, even in this case with only three monomers (in contrast to biological samples, where complexity is limited, facilitating database approaches). We refrain from drawing conclusions based on such an approach, as they are likely flawed. This – and our intentions to move beyond these simple systems – is the basis for our preferring tools developed for untargeted metabolomics (no specific product expectations), over the more obvious tools developed for proteomics.

2.3 Environment-Directed AA Condensation Experiments: Functional Examination

2.3.1 Reactivity testing using *p*NPA

In these experiments, the effects of product populations on the breakdown of para-nitrophenyl acetate (pNPA, colourless) to release para-nitrophenol (pNP, yellow) were observed, following this potentially very complex reaction system through the evolution of the yellow colour characteristic of free pNP (absorbance at 405 nm).




Figure S28. Breakdown of pNPA to yield pNP, turning the solution from colourless to yellow.

<u>Samples</u>: Unless otherwise stated, stock solutions were prepared from previously lyophilised product populations as outlined in Section 2.1, diluting the products of a reaction to a constant volume regardless of amounts of product produced. In some cases (where large amounts were yielded by all conditions tested), solutions of 0.5 mg/ml were also prepared (labelled as 'Constant Concentration, "CC", rather than 'Constant Volume', "CV").

<u>Assay</u>: A buffered substrate solution was prepared by adding 300 μ l of *p*-nitrophenyl acetate 0.1 M (in acetonitrile, for ease of handling) and 300 μ l of HEPES buffer 1 M to 11400 μ l of water. The final amount of acetonitrile present was 1.875%.

150 μ l of buffered substrate solution were then added to 50 μ l of product ensemble stock solution (giving a final substrate concentration of 1.875 mM). Kinetic measurements were performed in an Infinite M200 Pro Tecan plate reader (using the accompanying software for control and data capture) monitoring the absorbance of the *p*NP at 405 nm and at 25 °C, in a 96-well plate every 5 min for 2 h. At least 12 measurements were collected for each treatment (salt, mineral, mixing history).

<u>*Processing:*</u> Data was output from the instrument software in a spreadsheet format. Typical time-resolved traces can be observed in Figure S29. Initial rates were extracted using Microsoft Excel as the gradient (not constrained to the origin) of the plot of Abs₄₀₅ (in AU) against time (in seconds) over the first 60 minutes (close to linear in all cases).

Notes:

We note that while this is a common assay for esterase activity, catalysis of ester hydrolysis by the condensation products is not the only possible reaction type. We are interested in the effect of the complete ensemble of products on the reaction system and have made no attempt to identify the mechanism of pNP release (the complex set of competing pathways may include: ionic strength effects on uncatalysed reaction, inhibition of hydrolysis by recognition, disassembly of active catalytic assemblies on pNPA or pNP recognition, and other pathways).




Figure S29. Example plot of evolution of pNP (yellow colour, measured as Abs₄₀₅) over time, comparing product ensembles formed in the presence of NaCl or LiCl, and a Control lacking any products [error bars represent one standard deviation].




Figure S30 Results of reactivity testing using pNPA. Product ensembles produced with a variety of different environments are represented: (a) varying salts present, in (b) varying minerals present and (c) varying mixing history. Box plots (d), (e) and (f) comparing rates of pNP release from the same ensembles. Box plot (g) compares the rate of pNP release from ensembles produced from reactions with different mixing histories, diluted to a constant concentration of 0.5 mg/ml (rather than dissolving whatever products are yielded by a reaction to a fixed volume).

2.3.2 Recognition assay using ThT

This was carried out following an adaptation of an established approach.¹⁵ A stock solution of Thioflavin T (ThT) (Sigma) was prepared by dissolving 8 mg of ThT in 10 ml Tris buffer pH8 (Sigma), followed by filtration through a 0.2 μ m syringe filter. The working solution was prepared by diluting the stock into the buffer (1 ml stock to 50 ml buffer). 50 μ l of the peptide solution and 20 μ l of the ThT working solution were mixed in a 96 well-plate (Thermo Fisher). Fluorescence was measured after one hour of incubation using (Infinite® 200 PRO plate reader) by excitation at 444 nm and emission at 480 nm. Samples were measured in duplicate. Fluorescence values of the samples were compared to the ThT values (as a control).

2.3.3 Inspection of Assembly/Aggregation using TEM

Procedure

Carbon-coated copper grids (200 mesh) were glow discharged in air for 30 seconds. The support film was touched onto the peptide solution surface for 10 seconds, and excess solution was removed using filter paper. 20 μ l of negative stain (Nanovan; Nanoprobes) was applied and the mixture was blotted again using filter paper to remove any excess stain. The dried specimens were then imaged using an FEI Tecnai T20 Tranmission Electron Microscope (TEM) operating at 200 kV fitted with Gatan 794 Multiscan camera. Images were collected and converted to .tiff files using Gatan Microscopy Suite software.





Figure S31. Additional TEM images.

2.3.4 Observation of different properties of gels produced on addition of Ca²⁺ salts.

Following the difference in structural formation ability from amino acids' mixing history (Section 2.3.3), the difference of gelability was studied by peptides crosslinking with Ca²⁺. This was performed by mixing 500 μ l of each peptide solution (prepared in Section (2.1.4)) together with 2.5 μ l of 1 M CaCl₂, vortexing and leaving to stand overnight at room temperature. Gelability was verified by the inverted vial method (see Figure S32, in which those samples which are immobile were persistent in the position shown for periods > 1 h), and the products were observed using TEM following gelation, revealing dramatically different morphology (Figure S33).



 $\begin{array}{cccc} \mathsf{A}{+}\mathsf{V}{+}\mathsf{D} & \mathsf{D}{\Leftrightarrow}\,\mathsf{A}{\Leftrightarrow}\,\mathsf{V} & \mathsf{V}{\Leftrightarrow}\,\mathsf{A}{\Leftrightarrow}\,\mathsf{D} & \mathsf{A}{\Leftrightarrow}\,\mathsf{V}{\Leftrightarrow}\,\mathsf{D} \\ \mathsf{D}{\Leftrightarrow}\,\mathsf{V}{\Leftrightarrow}\,\mathsf{A} & \mathsf{V}{\Leftrightarrow}\,\mathsf{D}{\Rightarrow}\,\mathsf{A} & \mathsf{A}{\Leftrightarrow}\,\mathsf{D}{\Rightarrow}\,\mathsf{V} \end{array}$



Figure S32. Different ensembles produce materials with dramatically different degrees of gelation on addition of CaCl₂. Note: in image below, it is clear that in product ensembles without persistent gelling, clear solutions are observed rather than weaker gels.




Figure S33. TEM images showing assembly/aggregation of product ensembles on addition of Ca^{2+} salts (where outline is blue, scale bar = 0.2 μ m; where outline is green, scale bar = 0.5 μ m; where outline is black, scale bar = 2.0 μ m).

3 Environment-Directed Complex Mixture Condensation Experiments

3.1 Spark Discharge Mixture Preparation

Spark discharge mixtures were chosen as a complex mixture for this study, both as a classic 'intractable' complex mixture,¹⁶ and as they are known to contain a range of species (amino acids, hydroxyl acids, amines, etc.) amenable to simple condensation reactions. The spark discharge reaction product mixture ("SD Mix") used here as a model complex mixture was prepared from the combination of several SD reaction runs, using equipment similar to that of the 1950's Miller-Urey experiment.¹⁷ In each run, after careful cleaning and drying of the glassware, 400 mL of water (LC-MS grade) was added and the system sealed. The whole rig was pumped down three times to de-gas the water and finally after the third evacuation, the system was pressurised to 1 atm with gas mixture (40% methane, 40% ammonia and 20% hydrogen). Heating was applied to the main flask and, once boiling and recirculation was established, the 24 kV spark discharge was applied with a 10 sec alternating duty-cycle. Experiments were run for seven days, during which time the solution in the flask became deep brown in colour.

A total of ca. 0.5 L of product mixture was collected, combining the products of several runs. In order to produce a standardised mixture, free of large amounts of slowly-precipitating SiO₂ (dissolved from glassware), the mixture was then freeze-dried, redissolved in water (LC-MS grade) centrifuged (at 10k rpm for 1 h using a Beckman Coulter Avanti I-E centrifuge) and filtered (Millipore Durapore 0.22 μ m, HV type membranes), freeze-dried again and re-filtered (no observable residue). This yielded a light tan-coloured solution, containing approximately 1 mg/ml of soluble material.

3.2 Environment-Directed Complex Mixture Experiments: Synthesis

In this set of experiments, 4 ml of a standard SD mixture (see Section 3.1) was condensed (by dehydration) in the presence of a series of different minerals.

- 1. 4 ml of the standard SD mixture was added to each reaction vessel (open vial).
- 2. 0.2 g of a powdered mineral were added to each individual experiment. In addition, a control reaction with no mineral was set up (known as "Control (NM)"), and a control in which no condensation reaction took place (i.e. Step 3 was omitted and the SD mixture was stored at 4 °C) was set up (known as "Control (NR)").
- 3. A single dehydration step was performed in a fan-assisted oven at 115 °C for 24 h (a fixed arbitrary cycle time; all reactions performed together).
- 4. Reactions were then removed, and cooled to room temperature.
- 5. Each individual product mixture was dissolved in 4 ml of water, with sonication to aid dissolution of soluble species.
- Each individual product mixture solution was then filtered (0.2 μm, Pall Microsep centrifugal filter) to remove minerals and undissolved species.
 [n.b. centrifugal filters used to maximise and standardise product recovery]
- The filtrate of each reaction (and washings) was dialysed with a G2 Float-a-lyser (100-500 Da) cut-off (5 ml) for 24 h, to remove any small species and soluble salts.
- 8. Once the dialysis was completed, the samples (and washing, to avoid loss) were left to freeze-dry for 48 h.
- 9. The product mixtures were then redissolved in 0.5 ml water, and used/analysed without further treatment.

3.3 Environment-Directed Complex Mixture Experiments: Product Analysis

LC-MS analysis was accomplished in an adaptation of the general procedure described in Section 1.2, in which the linear gradient mixture of solvents A (water w/0.1% v/v formic acid) and B (acetonitrile w/0.1% v/v formic acid) was as follows over 40 min: 0 min – 0% B; 4 min – 0% B; 26 min – 100% B; 30 min – 100% B; 36 min – 0% B.

This LC-MS data was then processed and plotted as described in Section 2.2.1: peak picking and grouping, and gap-filling from raw data where no peaks were observed. PCA was performed on the resulting data as earlier (m/z and rt coordinates for each feature, with corresponding intensity for each sample), again with scaling. The results of this analysis are shown below, along with some sample EICs illustrating variance.




Figure S34 Plots of PCA analysis of results from condensation of SD mixture in the presence of different minerals. (*a*) Plot of first two PCs [in each case 'bubbles' represent 95% confidence limits & 'spots' represent individual measurements.] (*b*) Distribution of contributions to the first two principal components. (*c*) Plot of fraction of variance explained by these principal components.




Figure S35 (*a-c*) A range of plots (with different perspective) of PC-DFA analysis (using first 5 PCs) of results from condensation of SD mixture in the presence of different minerals; in each case 'spots' represent individual measurements & 'bubbles' represent two standard deviations around their mean. (d) Key to identify product ensembles, denoting mineral in whose presence they were produced, and matrix to clarify which ensembles overlap. Analysis conducted in R, calculated and plotted using rgl library.




Figure S36. Selected extracted ion chromatograms illustrating product distribution variance in products from experiments varying minerals present – Part 1 of 2. (a) m/z = 101.0715; (b) m/z = 102.0918; (c) m/z = 142.0507; (d) m/z = 166.0245; (e) m/z = 174.0582; (f) m/z = 176.0138; (g) m/z = 193.1387; (h) m/z = 203.1029; (i) m/z = 208.0464; (j) m/z = 218.1134. [lines = mean intensity from all measurements; shading around line represents one standard deviation around mean; intensities normalised relative to largest value in each plot]






Figure S37. Selected extracted ion chromatograms illustrating product distribution variance in products from experiments varying minerals present – Part 2 of 2. (a) m/z = 219.0614; (b) m/z = 223.1186; (c) m/z = 230.1610; (d) m/z = 242.0768; (e) m/z = 243.6842; (f) m/z = 244.1907; (g) m/z = 250.1538; (h) m/z = 252.0362; (i) m/z = 255.0590; (j) m/z = 262.0142; (k) m/z = 278.0520; (l) m/z = 300.7007; (m) m/z = 302.1963; (n) m/z = 303.2014; (o) m/z = 321.0014. [lines = mean intensity from all measurements; shading around line represents one standard deviation around mean; intensities normalised relative to largest value in each plot] Page 53





[i.e. $(Mean^{Mica} - Mean^{CtrlNM}) / Mean^{AllMin}$, where $Mean^{AllMin}$ is the mean intensity across all mineral environments; feature m/z and rt coordinates unlabelled, ordered by ascending m/z from left to right (as ca. 1800 features are plotted, lines are thin); Note: Fractional intensities can obscure smaller variations in intensity values below control]

m/z	RT	Alumina	Montmoril.	Mica	Goethite	Quartz	Natrolite	Silica	Control NM	Control NR
97.9695	1.34	4758	69677	33681	16645	16058	2196	5606	4939	5618
127.0504	1.81	68551	58297	43788	49177	39972	57185	46366	52738	51409
134.0449	1.75	99836	133759	190397	41431	222630	115551	194117	147357	150881
139.0616	3.53	152654	30072	144496	35296	111544	251713	55691	132252	255124
141.9590	1.34	4782	64603	30740	13840	13666	2288	5275	5135	5525
146.0449	1.78	87036	113150	34794	26925	44750	79386	55686	47217	109179
148.0606	1.56	120081	150471	175253	63832	155948	99580	146947	127985	140443
153.0296	2.24	227431	56357	121572	154220	93307	242439	89482	149671	150780
154.9075	1.33	51182	5449	85393	1454	281068	715202	117453	90043	111562
159.0767	2.18	589186	11389	263952	138981	294048	375657	266328	471051	407375
164.9208	1.41	5530	20731	2830	184940	423	438	2565	4133	2680
170.0564	2.24	225008	54874	113421	157426	87229	221947	87470	169678	148950
171.0401	2.24	488448	126381	265222	343306	206835	537263	199058	329210	342693
172.0352	3.51	129503	72064	54645	90577	39990	95659	52825	99017	76573
173.0564	1.82	76873	84218	54309	59553	48571	60996	56391	65510	65845
178.0618	7.41	2763628	6311	(034531	879851	663844	3073061	346802	1630504	1486776
179.0643	7.41	194134	712	93666	76034	63863	237517	32159	111906	132947
179.9027	1.33	42358	4730	67089	309	222620	589515	94597	72904	86542
180.0520	7.80	197320	486	108921	2630	45919	258156	39094	115844	128328
180.8954	1.42	20781	6367	18010	31555	83774	376297	24561	30865	39662
180.9018	1.33	50834	7588	80213	7304	272480	721551	114034	85450	106560
181.9036	1.33	75680	8830	122173	564	401927	1105939	171702	124189	160184
182.9025	1.33	495035	56202	831671	2357	1050730	1105335	1212096	872916	1100515
182.9490	1.42	48943	8708	65921	3760	256589	4500049	258123	184728	259250
192.0503	1.52	49502	64079	89095	10870	93252	50045	180044	78468	124572
192.0505	1.32	12819	66634	39021	746	72477	273	3946	1858	738
197.0569	1.99	7844	117767	6219	5885	5852	9261	20470	4817	12749
197.0509	7.81	408193	764	268779	75057	113953	613955	94487	361024	316185
202.1800	6.20	257497	86798	142268	253048	68661	191611	116563	168354	153304
202.1800	2.17	305726	21289	140575	70704	155592	193292	141139	247349	243870
	7.37	163523	3624	70417	52168	50560	47051	16703	67915	5852
211.0693 214.9171	1.33	29425	35492	38102	137076	48345	47651	50659	24568	26907
226.9517		103206	15894	25314	440	10436	21491	98459	131318	110926
230.8899	1.43	4070	59125	34762	29247	16110	934	5968	4360	4787
236.0672	1.33 9.45	160959	757	11177	302990	12653	2659	113890	72676	1550
306.8520	9.45	2793	7342	6861	103547	2500	1044	2275	1492	930
312.8616	1.42	39265	2765	42828	3713	172018	110177	69433	62845	58449
316.2121	1.55	177734	78867	98923	192092	49484	57021	74131	101567	78077
409.8053	1.33	7955	331	11710	307	78316	316181	30331	20724	27234
							1069344			len .
410.8048	1.33	24741	1366	37358	841	251392		96763	63423	86249 27493
432.2813	11.11	109273	42204	58621	119070	37839	94214	42197	76007	-
476.3068	11.37	237263	76414	119518	233922 350553	76433	182286	44337	150087	62969
520.3320	11.59	377373	107105	196390		125579	300242	71225	219531	116134
564.3592	11.80	465661	120584	238242	\$87860	161575	329568	89390	199009	156273
565.3618	11.80	120140	32883	65085	108573	44280	87258	25189	52615	43625
608.3844	11.99	368600	105699	220423	376512	157099	288826	81510	162958	156437
609.3892	11.99	104814	31378	62507	107352	46885	83717	24939	48329	47797
652.4113	12.16	230354	76527	150709	279921	124915	185245	57244	123853	119766
696.4368	12.32	119185	44948	83743	152768	75849	101000	29209	70794	61350

Figure S39. Table of selected features ordered by RT from the SD-mix experiment with different minerals. Features were selected from a full list based on absolute MS intensity (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

m/z	RT	Alumina	Montmoril.	Mica	Goethite	Quartz	Natrolite	Silica	Control NM	Control NR
214.9171	1.33	29425	35492	38102	137076	48345	4069	50659	24568	26907
312.8616	1.33	39265	2765	42828	3713	172018	110177	69433	62845	58449
410.8048	1.33	24741	1366	37358	841	251392	1069344	96763	63423	86249
409.8053	1.33	7955	331	11710	307	78316	316181	30331	20724	27234
230.8899	1.33	4070	59125	34762	29247	16110	934	5968	4360	4787
179.9027	1.33	42358	4730	67089	309	222620	589515	94597	72904	86542
182.9025	1.33	495035	56202	831671	2357	1050730	8359275	12:12:095	872916	1100515
154.9075	1.33	51182	5449	85393	1454	281068	715202	117453	90043	111562
180.9018	1.33	50834	7588	80213	7304	272480	721551	114034	85450	106560
181.9036	1.33	75680	8830	122173	564	401927	1105939	171702	124189	160184
97.9695	1.34	4758	69677	33681	16645	16058	2196	5606	4939	5618
141.9590	1.34	4782	64603	30740	13840	13666	2288	5275	5135	5525
194.9665	1.39	12819	66634	39021	746	72477	273	3946	1858	738
164.9208	1.41	5530	20731	2830	184940	423	438	2565	4133	2680
182.9490	1.42	48943	8708	65921	3760	256589	4500049	258123	184728	259250
180.8954	1.42	20781	6367	18010	31555	83774	376297	24561	30865	39662
306.8520	1.42	2793	7342	6861	103547	2500	1044	2275	1492	930
226.9517	1.43	103206	15894	25314	440	10436	21491	98459	131318	110926
192.0503	1.52	49502	64079	89095	10870	93252	50045	180044	78468	124572
148.0606	1.56	120081	150471	175253	63832	155948	99580	146947	127985	140443
1.8.9.0.0.000000000000000000000000000000	1.75	99836	133759	190397	41431	222630	115551	194117	147357	150881
134.0449								12	0	
146.0449	1.78	87036	113150	34794	26925	44750	79386	55686	47217	109179
127.0504	1.81	68551	58297	43788	49177	39972	57185	46366	52738	51409
173.0564	1.82	76873	84218	54309	59553	48571	60996	56391	65510	65845
197.0569	1.99	7844	117767	6219	5885	5852	9261	20470	4817	12749
205.0820	2.17	305726	21289	140575	70704	155592	193292	141139	247349	243870
159.0767	2.18	589186	11389	263952	138981	294048	375657	266328	471051	407375
170.0564	2.24	225008	54874	113421	157426	87229	221947	87470	169678	148950
171.0401	2.24	488448	126381	265222	343306	206835	537263	199058	329210	342693
153.0296	2.24	227431	56357	121572	154220	93307	242439	89482	149671	150780
172.0352	3.51	129503	72064	54645	90577	39990	95659	52825	99017	76573
139.0616	3.53	152654	30072	144496	35296	111544	251713	55691	132252	255124
202.1800	6.20	257497	86798	142268	253048	68661	191611	116563	168354	153304
211.0693	7.37	163523	3624	70417	52168	50560	47051	16703	67915	5852
179.0643	7.41	194134	712	93666	76034	63863	237517	32159	111906	132947
178.0618	7.41	263628	6311	1084531	879851	663844	3073061	346802	680504	1486776
180.0520	7.80	197320	486	108921	2630	45919	258156	39094	115844	128328
198.0622	7.81	408193	764	268779	75057	113953	613955	94487	361024	316185
236.0672	9.45	160959	757	11177	302990	12653	2659	113890	72676	1550
316.2121	10.80	177734	78867	98923	192092	49484	57021	74131	101567	78077
432.2813	11.11	109273	42204	58621	119070	37839	94214	42197	76007	27493
476.3068	11.37	237263	76414	119518	233922	76433	182286	44337	150087	62969
520.3320	11.59	377373	107105	196390	350553	125579	300242	71225	219531	116134
565.3618	11.80	120140	32883	65085	108573	44280	87258	25189	52615	43625
564.3592	11.80	465661	120584	238242	87860	161575	329568	89390	199009	156273
608.3844	11.99	368600	105699	220423	p76512	157099	288826	81510	162958	156437
609.3892	11.99	104814	31378	62507	107352	46885	83717	24939	48329	47797
652.4113	12.16	230354	76527	150709	279921	124915	185245	57244	123853	119766
696.4368	12.32	119185	44948	83743	152768	75849	101000	29209	70794	61350

Figure S40. Table of selected features ordered by RT from the SD-mix experiment with different minerals. Features were selected from a full list based on absolute MS intensity (appearing in top 20 for at least one condition); this is an arbitrary reduction of data for more detailed display, and it is important to note that no conclusion should be drawn on the significance of this selection due to the non-linear relationship between abundance and intensity. Intensities are averaged over experimental and analytical replicates.

3.4 Environment-Directed Complex Mixture Experiments: Functional Examination

3.4.1 Recognition assay using ThT

Procedure:

Following the same procedure as described in Section 2.3.2, with the modification that slightly smaller amounts of all materials were used (maintaining the same ratio: $25 \ \mu$ l 'sample' & 10 μ l ThT 'working solution'), and each sample tested in triplicate.

3.4.2 Inspection of Assembly/Aggregation using TEM

Procedure:

Following the same procedure described in Section 2.3.3, solutions of the product ensembles produced in Section 3.2 were inspected using TEM microscopy. Results are shown in Figures S41 & S42, and discussed below.

Observations:

- Observable morphological difference between populations: While in most samples more than one structure is present, observation of many images of the different populations at low magnification (Figure S41) shows that some populations are clearly distinct to others. (e.g. those produced in the presence of Goethite and Natrolite are clearly distinct). Observation at higher magnifications (Figure S42) reveals different structural detail in many samples, consistent with qualitative difference in the material present.
- Morphological observations not directly correlated with recognition assay results: Where populations have similar results in ThT recognition assay, they do not necessarily appear to produce morphologically similar assemblies when observed by TEM (e.g. those produced in the presence of Goethite and Natrolite are clearly distinct). This suggests that variation in recognition and assembly properties are not mediated by the same, simple, factor (e.g. amount of material present), but from qualitative differences between the products present.




Figure S41. TEM images showing assembly/aggregation of product ensembles from reaction of SD Mix in the presence of different minerals, at low magnification (scale bar = $2 \mu m$).




Figure S42. TEM images showing assembly/aggregation of product ensembles from reaction in the presence of different minerals, at low magnification (where outline is blue, scale bar = $0.2 \mu m$; *where outline is green, scale bar* = $0.5 \mu m$; *where outline is red, scale bar* = $1.0 \mu m$).

4 References

- 1. R: A language and environment for statistical computing. (R Foundation for Statistical Computing, Vienna, Austria., 2008). <u>http://www.r-project.org</u>
- 2. Chambers, M. C. *et al.* A cross-platform toolkit for mass spectrometry and proteomics. *Nat. Biotechnol.*, **30**, 918-920 (2012).
- 3. Smith, C. A., Want, E. J., O'Maille, G., Abagyan, R. & Siuzdak, G. XCMS: Processing mass spectrometry data for metabolite profiling using Nonlinear peak alignment, matching, and identification. *Anal. Chem.*, **78**, 779-787 (2006).
- 4. Long, W. in *Application Note 5991-5571EN*. (Agilent Technologies, Inc.). <u>http://www.agilent.com/cs/library/applications/5991-5571EN.pdf</u>
- 5. Commeyras, A. *et al.* Prebiotic synthesis of sequential peptides on the Hadean beach by a molecular engine working with nitrogen oxides as energy sources. *Polym. Int.*, **51**, 661-665 (2002).
- 6. Dick, J. M., LaRowe, D. E. & Helgeson, H. C. Temperature, pressure, and electrochemical constraints on protein speciation: Group additivity calculation of the standard molal thermodynamic properties of ionized unfolded proteins. *Biogeosciences*, **3**, 311-336 (2006).
- 7. Dick, J. M. Calculation of the relative metastabilities of proteins using the CHNOSZ software package. *Geochem. Trans.*, **9** (2008).
- 8. Anderson, G. M. *Thermodynamics of natural systems*. (Cambridge University Press, 2009).
- 9. James, G., Witten, D., Hastie, T. & Tibshirani, R. *An introduction to statistical learning: with applications in R.* Vol. 103 (Springer Science & Business Media, 2013).
- 10. Glansdorff, P. & Prigogine, I. Structure, stability and fluctuations. *New York, NY: Interscience* (1971).
- 11. Le, S., Josse, J. & Husson, F. FactoMineR: An R package for multivariate analysis. *J Stat Softw*, **25**, 1-18 (2008).
- 12. rgl: 3D Visualization Using OpenGL (2016). https://r-forge.r-project.org/projects/rgl/
- 13. Origin Pro 2016 (OriginLab, Northampton, MA, 2016). http://www.originlab.com
- 14. Venables, W. N., Ripley, B. D. & Venables, W. N. *Modern applied statistics with S.* 4th edn, (Springer, 2002).
- 15. Kyle, S., Felton, S. H., McPherson, M. J., Aggeli, A. & Ingham, E. Rational molecular design of complementary self-assembling peptide hydrogels. *Adv Healthc Mater*, **1**, 640-645 (2012).
- 16. Schwartz, A. W. Intractable mixtures and the origin of life. *Chem. Biodiversity*, **4**, 656-664 (2007).
- 17. Miller, S. L. A Production of amino acids under possible primitive earth conditions. *Science*, **117**, 528-529 (1953).