

## 4-Chloro-*N*-(4-cyano-2-nitrophenyl)-3-nitrobenzamide

Leroy Cronin,\*† Dave A. Adams, David J. Nightingale and James H. Clark

Department of Chemistry, University of York, Heslington, York YO10 5DD, England  
Correspondence e-mail: l.cronin@ed.ac.uk


Received 28 June 1999

Accepted 11 November 1999

The structure of the title compound,  $C_{14}H_7ClN_4O_5$ , comprises two nearly coplanar phenyl rings connected *via* an amido moiety.

### Comment

In the course of an investigation of the effect of water on the fluorodenitration of substituted benzonitriles, it was observed that hydrolysis of the  $-CN$  moiety was possible. In an attempted fluorination of 4-chloro-3-nitrobenzonitrile, the title compound, (I), was observed as a previously uncharacterized product in such reactions (Adams *et al.*, 1999).



In (I) (Fig. 1), the amido unit is planar with an O3—C8—N3—H3 torsion angle of 171 (2)°. However, the C10—C9—C8—O3 and C4—C5—N3—C8 torsion angles of 22.3 (4) and  $-28.0$  (4)°, respectively, indicate significant deviation from planarity with the amido group. In the chlorine-containing ring, the nitro group is twisted out of the plane of the phenyl ring [the O4—N4—C11—C10 torsion angle is 53.3 (4)°]. This is presumably due to the steric clash between the Cl atom and nitro group. In contrast, the nitro group *ortho* to the amido group on the other phenyl ring is more coplanar with the ring [the O2—N2—C6—C5 torsion angle is  $-14.5$  (4)°]. This conformation may be stabilized, at least in part, by a hydrogen-bond interaction between the N—H of the amido group and an O atom of the nitro group [N3...O2 2.662 (5), N3—H3...O2 2.06 (3) Å and N3—H3...O2 125 (3)°]. The planes of the phenyl rings are inclined to each other at an angle of 3.2 (3)°. Examination of the packing of the molecules

† Current address: Department of Chemistry, University of Edinburgh, West Mains Road, Edinburgh EH9 1NH, Scotland.




Figure 1

View of (I) showing the atom-numbering scheme and ellipsoids at the 50% probability level.

reveals a head-to-tail phenyl–phenyl interaction between adjacently stacked molecules of 3.77 (1) Å, indicating a weak intermolecular  $\pi$ -stacking interaction.

### Experimental

4-Chloro-3-nitrobenzonitrile (0.182 g, 1 mmol) was placed in a round-bottomed flask. Dimethyl sulfoxide (10 ml, distilled and stored under argon) was added and the solution heated to 353 K under argon. Tetramethylammonium fluoride (0.20 g, 1.70 mmol; TMAF $\cdot\frac{4}{3}$ H $_2$ O, prepared by drying the tetrahydrate under vacuum at 333 K for 2 d) was added to the solution. After 1 h, the reaction mixture was cooled in an ice bath. The organics were extracted into ether, washed well with water, dried (magnesium sulfate) and the ether removed on a rotary evaporator. The crude mixture was then recrystallized from acetonitrile to give yellow needles (Adams *et al.*, 1999).

#### Crystal data

$C_{14}H_7ClN_4O_5$   
 $M_r = 346.69$   
Triclinic,  $P\bar{1}$   
 $a = 7.804$  (13) Å  
 $b = 12.931$  (6) Å  
 $c = 7.49$  (4) Å  
 $\alpha = 91.59$  (16)°  
 $\beta = 112.9$  (2)°  
 $\gamma = 88.12$  (8)°  
 $V = 695$  (4) Å $^3$

$Z = 2$   
 $D_x = 1.656$  Mg m $^{-3}$   
Mo  $K\alpha$  radiation  
Cell parameters from 20 reflections  
 $\theta = 9$ –15°  
 $\mu = 0.312$  mm $^{-1}$   
 $T = 293$  (2) K  
Needle, yellow  
0.55 × 0.20 × 0.20 mm

#### Data collection

Rigaku four-circle AFC-6 diffractometer  
 $\omega$ – $2\theta$  scans  
Absorption correction: empirical via  $\psi$  scans (North *et al.*, 1968)  
 $T_{\min} = 0.782$ ,  $T_{\max} = 0.940$   
2453 measured reflections  
2453 independent reflections

1747 reflections with  $I > 2\sigma(I)$   
 $\theta_{\max} = 25.03$ °  
 $h = 0 \rightarrow 9$   
 $k = -15 \rightarrow 15$   
 $l = -8 \rightarrow 8$   
3 standard reflections every 150 reflections  
intensity variation: 0.5%

#### Refinement

Refinement on  $F^2$   
 $R[F^2 > 2\sigma(F^2)] = 0.046$   
 $wR(F^2) = 0.145$   
 $S = 1.053$   
2452 reflections  
220 parameters  
H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0735P)^2 + 0.1533P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.24$  e Å $^{-3}$   
 $\Delta\rho_{\min} = -0.31$  e Å $^{-3}$

**Table 1**

Selected geometric parameters (Å, °).

C11–C12	1.724 (3)	O5–N4	1.219 (6)
O1–N2	1.225 (4)	N1–C1	1.147 (4)
O2–N2	1.234 (3)	N2–C6	1.465 (4)
N3–C8	1.377 (5)	O3–C8	1.210 (4)
N3–C5	1.409 (4)	N4–C11	1.472 (4)
O4–N4	1.213 (5)	C1–C2	1.445 (4)
C8–N3–C5	125.4 (3)	O5–N4–C11	117.7 (3)
O1–N2–O2	122.5 (2)	N1–C1–C2	177.8 (3)
O1–N2–C6	118.3 (2)	O3–C8–N3	123.9 (3)
O2–N2–C6	119.2 (2)	O3–C8–C9	121.1 (3)
O4–N4–O5	124.9 (3)	N3–C8–C9	114.9 (3)
O4–N4–C11	117.3 (3)		

The H atom on N3 was located in a difference Fourier synthesis. It was allowed to refine positionally with  $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{N3})$ . The phenyl H atoms were placed geometrically and thereafter refined using a riding model with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

Data collection: *TEXSAN* (Molecular Structure Corporation, 1993); cell refinement: *TEXSAN*; data reduction: *TEXSAN*; program(s) used to solve structure: *SHELXS86* (Sheldrick, 1985);

program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993); molecular graphics: *ORTEP* (Johnson, 1965); software used to prepare material for publication: *SHELXL93*.

We thank Victrex plc for financial support (to DA), the EPSRC Clean Technology Unit for a studentship (to DN) and the Royal Academy of Engineering/EPSRC for a Clean Technology Fellowship (to JHC).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: BM1358). Services for accessing these data are described at the back of the journal.

## References

- Adams, D. A., Clark, J. H., McFarland, H. & Nightingale, D. J. (1999). *J. Fluorine Chem.* **94**, 51–54.
- Johnson, C. K. (1965). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee, USA.
- Molecular Structure Corporation (1993). *TEXSAN*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (1985). *SHELXS86*. *Crystallographic Computing 3*, edited by G. M. Sheldrick, C. Krüger & R. Goddard, pp. 175–189. Oxford University Press.
- Sheldrick, G. M. (1993). *SHELXL93*. University of Göttingen, Germany.